

02-02-06

A

JCS44 U.S. PTO
01/31/00

JCS30 U.S. PTO
09/495823
01/31/00

UTILITY PATENT APPLICATION TRANSMITTAL (Only for new nonprovisional applications under 37 CFR 1.53(b))	Attorney Docket No. 5800-79
	First Inventor or Application Identifier: Glucksmann et al.
	Title of Invention: 22438, 23553, 25278, AND 26212 NOVEL HUMAN SULFATASES
	Express Mail Label No. EL039496501US

ADDRESS TO: ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
WASHINGTON, DC 20231

Transmitted herewith for filing in the United States Patent Office is a patent application for:

Inventors: Maria Alexandra Glucksmann
Mark Williamson

1. ☒ The Filing Fee has been calculated as shown below:
(Submit an original, and a duplicate for fee processing)

	No. Filed	No. Extra	Small Entity		Large Entity	
			Rate	Fee 0	Rate	Fee 0
BASIC FEE				\$0		\$ 690
TOTAL CLAIMS:	954 - 20 =	934		X 9 = \$0		x 18 = \$16,812
INDEP CLAIMS:	11 - 3 =	8		X 39 = \$0		x 78 = \$ 624
<input checked="" type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED				+130 = \$		+260 = \$ 260
*If the difference in Column 1 is less than zero, enter "0" in Column 2.				TOTAL \$		TOTAL \$ 18,386

The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 16-0605.

- a. ☐ Fees required under 37 CFR 1.16 (National filing fees).
b. ☒ Fees required under 37 CFR 1.17 (National application processing fees).
☐ A check in the amount of \$ ___ for the filing fee is enclosed.
☒ The above filing fee will be paid along with Applicant(s) Response to the Notice to File Missing Parts.
2. ☒ Specification; Total Pages 119
3. ☒ 25 Sheets of Formal Drawing(s) (35 USC 113)

4. ☐ Declaration and Power of Attorney; [Total Pages ____]
- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 16 completed)
- i. ☐ DELETION OF INVENTOR(S) Signed statement
attached deleting inventor(s) named in the prior
application, see 37 CFR 1.63(d)(2) & 1.33(b).
5. ☐ Microfiche Computer Program (Appendix)
6. ☐ Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☐ Computer Readable Copy
- b. ☐ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment Papers (cover sheet & document(s) (including a check for
the \$40.00 fee)
8. ☐ 37 CFR 3.73(b) Statement (when there is an assignee); ☐ Power of Attorney
9. ☐ English Translation Document (if applicable)
10. ☐ Information Disclosure Statement (IDS)/PTO-1449; ____ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
13. ☐ Small Entity Statement(s)
☐ Statement filed in prior application; status still proper and desired.
14. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
Foreign Priority is
15. ☐ Other: ____

16. **If a CONTINUING APPLICATION**, check appropriate box and supply the requisite
information below and in a preliminary amendment:

☐ Continuation ☐ Divisional ☐ Continuation in Part (CIP)
of prior Application No: ____; Filed ____

Prior Application Information: Examiner

Group/Art Unit:

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is
supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby
incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted
application parts.

17. **CORRESPONDENCE ADDRESS**

Customer Number or Bar Code Label **000826**

Attention Of: Anne Brown

Signature: _____

Anne Brown

Attorney of Record: Anne Brown

Attorney Registration No. 36,463

Tel Raleigh Office (919) 420-2200

Fax Raleigh Office (919) 420-2260

ALSTON & BIRD LLP

P.O. Drawer 34009

Charlotte NC 28234-4009

"Express Mail" mailing label number EL039496501US

Date of Deposit January 31, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner For Patents, Washington, DC 20231.

Nora C Martinez

Nora C. Martinez

RTA01/2073097v1

Attorney's Docket No. 5800-79



#3
A122-0
PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE *A*

In re: Glucksmann *et al.*
Appl. No.: 09/495,823
Filed: January 31, 2000
For: 22438, 23553, 25278, and 26212 Novel Human Sulfatases

June 5, 2000

Box Missing Parts
Assistant Commissioner for Patents
Washington, DC 20231

PRELIMINARY AMENDMENT

Dear Sir:

Please amend the above-identified application as follows:

In The Claims:

✓ Please cancel claims 1-242, without prejudice to or disclaimer of the subject matter contained therein.

Please add the following new claims:

A1
--243. An isolated nucleic acid molecule selected from the group consisting of:

- a) a nucleic acid molecule comprising a nucleotide sequence which is at least 45% identical to the nucleotide sequence of SEQ ID NO:2, 4, 6, or 8, the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____, or a complement thereof;
- b) a nucleic acid molecule comprising a fragment of at least 15 nucleotides of the nucleotide sequence of SEQ ID NO:2, 4, 6, or 8, the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____,

_____, _____, or _____, or a complement thereof;

c) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____;

d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____, wherein the fragment comprises at least 12 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____; and

e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:2, 4, 6, or 8, or a complement thereof under stringent conditions.

244. The isolated nucleic acid molecule of claim 243, which is selected from the group consisting of:

a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:2, 4, 6, or 8, the cDNA insert of any one the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____, or a complement thereof; and

b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____.

245. The nucleic acid molecule of claim 243 further comprising vector nucleic acid sequences.

246. The nucleic acid molecule of claim 243 further comprising nucleic acid sequences encoding a heterologous polypeptide.

247. A host cell which contains the nucleic acid molecule of claim 243.

248. The host cell of claim 247 which is a mammalian host cell.

249. A nonhuman mammalian host cell containing the nucleic acid molecule of claim 243.

250. An isolated polypeptide selected from the group consisting of:

a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____;

- b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:2, 4, 6, or 8, or a complement thereof under stringent conditions; and
- c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 45% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:2, 4, 6, or 8, or a complement thereof.

251. The isolated polypeptide of claim 250 comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____.

252. The polypeptide of claim 250 further comprising heterologous amino acid sequences.

253. An antibody which selectively binds to a polypeptide of claim 250.

254. A method for producing a polypeptide selected from the group consisting of:
- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____, ____, ____, or ____;
- b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers ____.

_____, _____, or _____, wherein the fragment comprises at least 12 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____; and

c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Numbers _____, _____, _____, or _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:2, 4, 6, or 8, or a complement thereof under stringent conditions; comprising culturing the host cell of claim 247 under conditions in which the nucleic acid molecule is expressed.

255. The method of claim 252 wherein said polypeptide comprises the amino acid sequence of SEQ ID NO:1, 3, 5, or 7.

256. A method for detecting the presence of a polypeptide of claim 250 in a sample, comprising:

- a) contacting the sample with a compound which selectively binds to a polypeptide of claim 250; and
- b) determining whether the compound binds to the polypeptide in the sample.

257. The method of claim 256, wherein the compound which binds to the polypeptide is an antibody.

258. A kit comprising a compound which selectively binds to a polypeptide of claim 250 and instructions for use.

259. A method for detecting the presence of a nucleic acid molecule of claim 243 in a sample, comprising the steps of:

- a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and
- b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample.

260. The method of claim 259, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

261. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 243 and instructions for use.

262. A method for identifying a compound which binds to a polypeptide of claim 250 comprising the steps of:

- a) contacting a polypeptide, or a cell expressing a polypeptide of claim 250 with a test compound; and
- b) determining whether the polypeptide binds to the test compound.

263. The method of claim 262, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detecting of test compound/polypeptide binding;
- b) detection of binding using a competition binding assay;
- c) detection of binding using an assay for GPCR-like-mediated signal transduction.

264. A method for modulating the activity of a polypeptide of claim 250 comprising contacting a polypeptide or a cell expressing a polypeptide of claim 250 with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

265. A method for identifying a compound which modulates the activity of a polypeptide of claim 250, comprising:

- a) contacting a polypeptide of claim 250 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

266. A method for identifying an agent that modulates the level of expression of a nucleic acid molecule of claim 243 in a cell, said method comprising contacting said agent with the cell expressing said nucleic acid molecule such that said level of expression of said nucleic acid molecule can be modulated in said cell by said agent and measuring said level of expression of said nucleic acid molecule.

267. A method for modulating the level of expression of a nucleic acid molecule of claim 243, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level of expression of the nucleic acid molecule.

268. A pharmaceutical composition containing any of the polypeptides in claim 250 in a pharmaceutically acceptable carrier.--

In re: Glucksmann *et al.*
Appl. No.: 09/495,823
Filed: January 31, 2000
Page 8



REMARKS

The foregoing amendments to the claims are fully supported in the specification as originally filed. Accordingly, the foregoing amendments to the claims do not add new matter; their entry is therefore respectfully requested.

Applicants believe that the present application is now in condition for examination. If the Examiner believes, for any reason, that personal communication will expedite prosecution of this application, the Examiner is invited to telephone the undersigned at the number provided.

Prompt and favorable consideration of the foregoing amendments, and entry of the same into the present application, are respectfully requested.

It is not believed that extensions of time or fees for net addition of claims are required, beyond those that may otherwise be provided for in documents accompanying this paper. However, in the event that additional extensions of time are necessary to allow consideration of this paper, such extensions are hereby petitioned under 37 CFR § 1.136(a), and any fee required therefore (including fees for net addition of claims) is hereby authorized to be charged to Deposit Account No. 16-0605.

Respectfully submitted,

W. Murray Spruill
Registration No. 32,943

ALSTON & BIRD LLP

P.O. Drawer 34009
Charlotte, NC 28234
Tel Raleigh Office (919) 420-2200
Fax Raleigh Office (919) 420-2260

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Assistant Commissioner For Patents, Washington, DC 20231, on June 5, 2000.

Nancy Cochran

22438, 23553, 25278, AND 26212 NOVEL HUMAN SULFATASES

FIELD OF THE INVENTION

5 The present invention relates to newly identified human sulfatases. In particular, the invention relates to sulfatase polypeptides and polynucleotides, methods of detecting the sulfatase polypeptides and polynucleotides, and methods of diagnosing and treating sulfatase-related disorders. Also provided are vectors, host cells, and recombinant methods for making and using the novel molecules.

BACKGROUND OF THE INVENTION

10

The biology and functions of the reversible sulfation pathway catalyzed by human sulfotransferases and sulfatases has been reviewed by Coughtrie *et al.* (*Chemico-Biological Interactions* 109: 3-27 (191998)). This review, summarized below, focuses on the sulfation of small molecules carried out by cytosolic sulfotransferases rather than the sulfation of macromolecules and lipids catalyzed by membrane-associated sulfotransferases.

15

Sulfation functions in the metabolism of xenobiotic compounds, steroid biosynthesis, and modulating the biological activity and inactivation and elimination of potent endogenous chemicals such as thyroid hormones, steroids and catechols. This pathway is reversible, comprising the sulfotransferase enzymes that cause the sulfation and the sulfatases that hydrolyze the sulfate esters formed by the action of the sulfotransferases. Accordingly, the interplay between these families regulates the availability and biological activity of xenobiotic and endogenous chemicals. The sulfatases, including the arylsulfatases (ARS), are located in lysosomes or endoplasmic reticulum.

20

25

The presence of sulfated components depends upon the availability of key members of the sulfate pathway, i.e., substrate and activated sulfate donor molecule (co-substrate) and the balance between sulfation and sulfate conjugate hydrolysis that depends upon the activity and localization of the sulfotransferases and the sulfatases.

5 Essentially, divalent sulfate is converted to adenosine 5' phosphosulfate (PAPS) by hydrolysis of ATP. This compound is in turn converted to 3' phosphoadenosine 5' phosphosulfate by hydrolysis of ATP to ADP. This compound is then converted to adenosine 3' 5' biphosphate concurrently with the formation of 4-nitrophenolsulfate from 4-nitrophenol. An ARS would then cleave the monovalent sulfate from the 4-
10 nitrophenolsulfate to produce the original 4-nitrophenol. This forms the basis for the sulfation system in humans. Over- or under-production of any of these key molecules can result in sulfate-related disorders. For example, the brachymorphic mouse has a connective tissue disorder that results from a defect in PAPS formation that causes undersulfated cartilage proteoglycans.

15 ARS enzymes and their genes have been associated with specific genetic diseases. ARSA is located in the lysosomes and removes sulfate from sulfated glycolipids. A deficiency of ARSA has been associated with metachromatic leukodystrophy and multiple sulfatase deficiency (MSD). ARSB is located in lysosomes and has, as an endogenous substrate, dermatan sulfate and chondroitin sulfate. A
20 deficiency of ARSB is associated with Maroteaux-Lamy syndrome and MSD. ARSC is located in the endoplasmic reticulum and has, as its endogenous substrate, cholesterol sulfate and steroid sulfates. A deficiency of ARSC is associated with X-linked ichthyosis and MSD. ARSD may be associated with MSD. ARSE has been associated with chondrodysplasia punctata and MSD. ARSF may be associated with MSD. ARSC
25 hydrolyses sulfate esters on a wide range of steroids and cholesterol. ARSs also hydrolyse sulfate conjugates of xenobiotics.

MSD results from an inability to perform a co- or post-translational modification of a cysteine residue to serine semialdehyde (2-oxo-3-propionic acid). This residue is conserved in all eukaryotic sulfatases described by Coughtrie *et al.* ARSC may have a
30 very broad specificity, extending to iodothyronine sulfates and a number of sulfate conjugates of xenobiotic phenols.

The kinetic and catalytic properties of ARS enzymes in isolation, important for understanding substrate specificity and the physical and chemical properties of enzymes and substrates that allow substrate preference, have been characterized recently based on recombinant enzyme systems. For the expression of the human sulfotransferases, COS and V79 cells have been used. Coughtrie *et al.* have constructed and characterized V79 cell lines stably expressing ARSA, ARSB, and ARSC. These cell lines exhibited the expected substrate preferences of the three enzymes among the substrates 4-nitrocatechol sulfate, estrone sulfate, and dehydroepiandrosterone sulfate (DHEAS).

The sulfation of small molecules can be broadly divided into the areas of chemical defense, hormone biosynthesis, and bioactivation. It was originally viewed that sulfation protected against the toxic effects of xenobiotics in that sulfate conjugates are more readily excreted in urine or bile and generally exhibit reduced pharmacological/biological activity relative to the parent compound. Many drugs and other xenobiotics are conjugated with sulfate. Many phenolic metabolites of the cytochrome P450 mono-oxygenase system are excreted as sulfate conjugates.

Further, potent endogenous chemicals, such as steroids and catecholamines are found at high levels as circulating sulfate conjugates. For example, greater than 90% of circulating dopamine exists as the sulfated form. Sulfation is also suggested to play a role in the inactivation of potent steroids such as estrogens and androgens. Accordingly, sulfation is important in metabolism and homeostasis of such compounds in humans.

DHEAS is the major circulating steroid in humans and estrone sulfate is the major estrogen. These chemicals act as precursors of estrogens and androgens. Extremely large quantities of such steroids or estrogens may occur during various stages of development, such as pregnancy. Estrone sulfate is a precursor for β -estradiol synthesis. In breast cancer cells it is hydrolysed by steroid sulfatase (ARSC) to estrone which is then converted to β -estradiol by action of another enzyme. Accordingly, ARSC is important for maintaining active estrogen. It is thus an important therapeutic target for the treatment of breast cancer.

Cholesterol sulfate, synthesized in the skin epidermis, may have a role in keratinocyte differentiation. Accordingly, hydrolysis of cholesterol sulfate by steroid sulfatase may be important in skin formation and differentiation. This is the major organ affected in X-linked ichthyosis caused by mutations in ARSC.

Although sulfation may widely serve to detoxify potent compounds, some sulfate conjugates are more biologically active than the corresponding parent compound.

Minoxidil and cicletanine are activated upon sulfation. Further, an inhibitor of ARSC was shown to potentiate the memory enhancing effect of DHEAS. This suggests a role

5 for sulfates and sulfation in the central nervous system.

An important example of bioactivation by means of sulfation, however, occurs with dietary and environmental mutagens and carcinogens. For a large number of these, sulfation is the terminal step in the pathway to metabolic activation. Examples of such chemicals include aromatic amines (including heterocyclic amines) and benzylic

10 alcohols of chemicals such as polycyclic aromatic hydrocarbons, safrole, and estragole.

The sulfatase gene family has been reviewed in Parenti *et al.* (*Current Opinion in Genetics and Development* 7:386-391 (1997)), summarized below.

The sulfatase family of enzymes is functionally and structurally similar.

Nevertheless, these enzymes catalyze the hydrolysis of sulfate ester bonds from a wide variety of substrates ranging from complex molecules such as glycosaminoglycans and sulfolipids to steroid sulfates (see also Coughtrie *et al.*, above). Several human genetic disorders result from the accumulation of intermediate sulfate compounds that result from a deficiency of single or multiple sulfatase activities. A subset of sulfatase, ARS, is characterized by the ability to hydrolyze sulfate esters of chromogenic or fluorogenic

15 aromatic compounds such as *p*-nitrocatechol sulfate and 4-methylumbelliferyl sulfate.

Desulfation is required to degrade glycosaminoglycans, heparan sulfate, chondroitin sulfate and dermatan sulfate and sulfolipids. Steroid sulfatase differs from other members of the family with respect to subcellular localization. It is localized in the microsomes rather than in lysosomes. Further, ARSD, ARSE, and ARSF are also non-lysosomal, being localized in the endoplasmic reticulum or Golgi compartment.

25 The natural substrate of ARSA is cerebroside sulfate. Associated diseases are MLD and MSD. The natural substrate of ARSB is dermatan sulfate. The disease associated with this enzyme is MPSVI and MSD. The natural substrate of ARSC/STS is sulfated steroids. Diseases associated with this enzyme are XLI and MSD. The natural substrates of ARSD-F are unknown. The natural substrates of iduronate-2-sulfate sulfatase (IDS) are dermatan sulfate and herparan sulfate. Diseases associated with this enzyme are MPSII and MSD. The natural substrate of galactose 6-sulfatase is keratan

sulfate and chondroitin 6-sulfate. Diseases associated with this enzyme include MPSIVA and MSD. The natural substrate of glucosamine-6-sulfatase is heparan sulfate and keratan sulfate. A disease associated with this enzyme is MPSIIID and MSD. The natural substrate of glucuronate-2-sulfatase is heparan sulfate. The natural substrate of glucosamine-3-sulfatase is heparan sulfate.

Sulfatases are activated through conversion of a cysteine residue as described above. The conversion is required for catalytic activity and is defective in MSD. It is likely that all sulfatases undergo the same modification. The substitution of this cysteine was shown to destroy the enzymatic activity of N-acetyl galactosamine-4-sulfatase (ARSB). It has been shown that the modified residue and a metal ion are located at the base of a substrate binding pocket.

Nine human sulfatase genes are known and murine rat, goat, or avian orthologs for some of these have been identified. A high degree of similarity occurs particularly in the amino terminal region which contains accordingly a potential consensus sulfatase signature.

Sulfatases, as discussed above, are associated with human disease. Most sulfatase deficiencies cause lysosomal storage disorders. The mucopolysaccharidoses contain various associations of mental retardation, facial dysmorphisms, skeletal deformities, hepatosplenomegaly, and deformities of soft tissues caused by deficiencies of sulfatases acting on glycosaminoglycans. In metachromatic leukodystrophy, a deficiency of ARSA causes the storage of sulfolipids in the central and peripheral nervous systems, leading to neurologic deterioration. X-linked ichthyosis is caused by STS deficiency leading to increased cholesterol sulfate levels. MSD, a disorder in which all sulfatase activities are simultaneously defective, was shown to result from a defect in the co- or post-translational processing of sulfatases.

Accordingly, sulfatases are a major target for drug action and development. Therefore, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown sulfatases. The present invention advances the state of the art by providing previously unidentified human sulfatases.

SUMMARY OF THE INVENTION

Novel sulfatase nucleotide sequences, and the deduced sulfatase polypeptides
5 are described herein. Accordingly, the invention provides isolated sulfatase nucleic acid
molecules having the sequences shown in SEQ ID NOS:2, 4, 6, and 8 or in the cDNA
deposited as ATCC Nos. _____ on _____, respectively ("the deposited cDNA"),
and variants and fragments thereof.

It is also an object of the invention to provide nucleic acid molecules encoding
10 the sulfatase polypeptides, and variants and fragments thereof. Such nucleic acid
molecules are useful as targets and reagents in sulfatase expression assays, are applicable
to treatment and diagnosis of sulfatase-related disorders and are useful for producing
novel sulfatase polypeptides by recombinant methods.

The invention thus further provides nucleic acid constructs comprising the
15 nucleic acid molecules described herein. In a preferred embodiment, the nucleic acid
molecules of the invention are operatively linked to a regulatory sequence. The
invention also provides vectors and host cells for expressing the sulfatase nucleic acid
molecules and polypeptides, and particularly recombinant vectors and host cells.

In another aspect, it is an object of the invention to provide isolated sulfatase
20 polypeptides and fragments and variants thereof, including a polypeptide having the
amino acid sequence shown in SEQ ID NOS:1, 3, 5 or 7 or the amino acid sequences
encoded by the deposited cDNAs. The disclosed sulfatase polypeptides are useful as
reagents or targets in sulfatase assays and are applicable to treatment and diagnosis of
sulfatase-related disorders.

The invention also provides assays for determining the activity of or the presence
25 or absence of the sulfatase polypeptides or nucleic acid molecules in a biological sample,
including for disease diagnosis. In addition, the invention provides assays for
determining the presence of a mutation in the polypeptides or nucleic acid molecules,
including for disease diagnosis.

30 A further object of the invention is to provide compounds that modulate
expression of the sulfatase for treatment and diagnosis of sulfatase-related disorders.

Such compounds may be used to treat conditions related to aberrant activity or expression of the sulfatase polypeptides or nucleic acids.

The disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment of sulfatase related disorders. The compositions include sulfatase polypeptides, nucleic acids, vectors, transformed cells and related variants thereof. In particular, the invention relates to the diagnosis and treatment of sulfatase-related disorders including, but not limited to disorders as described in the background above, further herein, or involving a tissue shown in the figures herein.

In yet another aspect, the invention provides antibodies or antigen-binding fragments thereof that selectively bind the sulfatase polypeptides and fragments. Such antibodies and antigen binding fragments have use in the detection of the sulfatase polypeptide, and in the prevention, diagnosis and treatment of sulfatase related disorders.

The sulfatases disclosed herein are designated as follows: 22438, 23553, 25278, and 26212.

DESCRIPTION OF THE DRAWINGS

Figure 1 shows the 22438 sulfatase cDNA sequence (SEQ ID NO:2) and the deduced amino acid sequence (SEQ ID NO:1).

Figure 2 shows a 22438 sulfatase hydrophobicity plot.

Figure 3 shows an analysis of the 22438 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

Figure 4 shows an analysis of the 22438 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the

first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, an amidation site is found from about amino acids 56-59, an EGF-like domain cysteine pattern signature found from about amino acids 260-271, and a sulfatase signature is found from about amino acids 129-138.

Figure 5 shows the 23553 sulfatase cDNA sequence (SEQ ID NO:4) and the deduced amino acid sequence (SEQ ID NO:3).

Figure 6 shows a 23553 sulfatase hydrophobicity plot.

Figure 7 shows an analysis of the 23553 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

Figure 8 shows an analysis of the 23553 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual modified residue is the last amino acid residue. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, a sulfatase signature is found from about amino acids 85-97.

Figure 9 shows relative expression of the 23553 sulfatase mRNA in normal and cancerous human tissues.

Figure 10 shows the 25278 sulfatase cDNA sequence (SEQ ID NO:6) and the deduced amino acid sequence (SEQ ID NO:5).

Figure 11 shows a 25278 sulfatase hydrophobicity plot.

Figure 12 shows an analysis of the 25278 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

5 Figure 13 shows an analysis of the 25278 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual modified residue is the last amino acid residue. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, amidation sites are found from about amino acids 312-315 and 541-544, and sulfatase signatures are found from
10 about amino acids 139-148 and 91-103.
15

Figure 14 shows relative expression of sulfatase mRNA in normal and cancerous human tissues.

20 Figure 15 shows the 26212 sulfatase cDNA sequence (SEQ ID NO:8) and the deduced amino acid sequence (SEQ ID NO:7).

Figure 16 shows a 26212 sulfatase hydrophobicity plot.

25 Figure 17 shows an analysis of the 26212 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

30 Figure 18 shows an analysis of the 26212 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino

acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual modified residue is the last amino acid residue. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, sulfatase signature sites are found from about amino acids 168-177 and 120-132.

DETAILED DESCRIPTION OF THE INVENTION

10 Sulfatase Polypeptides

The invention is based on the identification of the novel human 22438 sulfatase. *In situ* hybridization experiments showed that this sulfatase is expressed in the following monkey tissues: sub-populations of DRG neurons (mainly in small and medium sized neurons), in spinal cord (interneurons and motor neurons), and in the brain. The sulfatase is also expressed in human brain. The sulfatase cDNA was identified based on consensus motifs or protein domains characteristic of sulfatases and, in particular, arylsulfatase. BLAST analysis has shown homology with human arylsulfatase E, a human iduronate-2-sulfatase, human N-acetylgalactosamine-6-sulfatase, murine arylsulfatase A, and human arylsulfatase A. However, some homology has also been found with other arylsulfatases from various mammalian species, including, but not limited to, human arylsulfatase D, E, F, and B.

The invention is also based on the identification of the novel human 23553 sulfatase. Taqman analysis has shown positive differential expression in breast and colon cancer and in colonic metastases to the liver (Figure 9). This sulfatase has been identified as a glucosamine-6-sulfatase based on ProDom matches and BLAST analysis. Some homology has also been found to human arylsulfatase A, human N-acetylglucosamine-6-sulfatase, and human iduronate-2-sulfatase.

The invention is also based on the identification of the novel human 25278 sulfatase. The sulfatase is differentially expressed in human colon cancer and in colonic metastases to the liver, as determined by Taqman analysis. This sulfatase has been identified as a N-acetylgalactosamine-4-sulfatase by ProDom matching and

BLAST homology alignment. Further, based on BLAST analysis, some homology has also been shown to arylsulfatase B and arylsulfatase A.

5 The invention is also based on the identification of the novel human 26212 sulfatase. This sulfatase has been identified as an arylsulfatase by ProDom matching and BLAST sequence alignment. Homology has been shown to arylsulfatase B. Some homology has also been found with arylsulfatase F, E, D, and A, as well as with iduronate 2 sulfatase. Arylsulfatase B is also known as N-acetylgalactosamine-4-sulfatase.

10 Specifically, newly-identified human genes, termed 22438, 23553, 25278, and 26212 sulfatases are provided. These sequences, and other nucleotide sequences encoding the sulfatase proteins or fragments and variants thereof, are referred to as "22438, 23553, 25278, and 26212 sulfatase sequences."

15 Plasmids containing the sulfatase cDNA inserts were deposited with the Patent Depository of the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Virginia, on _____, and assigned Patent Deposit Numbers _____, respectively. The deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. The deposits were made merely as a convenience for those of skill in the art and is not an admission that a
20 deposit is required under 35 U.S.C. § 112.

The sulfatase cDNA was identified in human cDNA libraries. Specifically, expressed sequence tags (EST) found in human cDNA libraries, were selected based on homology to known sulfatase sequences. Based on such EST sequences, primers were designed to identify a full length clone from a human cDNA library. Positive clones
25 were sequenced and the overlapping fragments were assembled. The 22438, 23553, 25278, and 26212 sulfatase amino acid sequences are shown in Figures 1, 5, 10, and 15, respectively, and SEQ ID NOS:1, 3, 5, and 7. The 22438, 23553, 25278, and 26212 sulfatase cDNA sequences are shown in Figures 1, 5, 10, and 15 and SEQ ID NOS:2, 4, 6, and 8.

30 Analysis of the assembled sequences revealed that the cloned cDNA molecules encoded sulfatase-like polypeptides. BLAST analysis indicated that the 23553 sulfatase is a glucosamine-6-sulfatase, that the 25278 sulfatase is an N-

acetylgalactosamine-4-sulfatase, that the 22438 is an arylsulfatase with highest homology to arylsulfatase A and E genes and that the 26212 sulfatase is an arylsulfatase with highest homology to the arylsulfatase B gene (N-acetylgalactosamine-4-sulfatase).

5 The sulfatase sequences of the invention belong to the sulfatase family of molecules having conserved functional features. The term "family" when referring to the proteins and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having sufficient amino acid or nucleotide sequence identity as defined herein to provide a specific function. Such family
10 members can be naturally-occurring and can be from either the same or different species. For example, a family can contain a first protein of murine origin and an ortholog of that protein of human origin, as well as a second, distinct protein of human origin and a murine ortholog of that protein.

 The 22438 sulfatase gene encodes an approximately 2175 nucleotide mRNA
15 transcript having the corresponding cDNA set forth in SEQ ID NO:2. This transcript has an open reading frame which encodes a 525 amino acid protein (SEQ ID NO:1).

 The 23553 sulfatase gene encodes an approximately 4321 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:4. This transcript has an open reading frame which encodes an 871 amino acid protein (SEQ ID NO:3).

20 The 25278 sulfatase gene encodes an approximately 2940 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:6. This transcript has an open reading frame which encodes a 569 amino acid protein (SEQ ID NO:5).

 The 26212 sulfatase gene encodes an approximately 2266 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:8. This transcript has
25 an open reading frame which encodes a 550 amino acid protein (SEQ ID NO:7).

 Prosite program analysis was used to predict various sites within the 22438 sulfatase protein as shown in Figure 4.

 Prosite program analysis was used to predict various sites within the 23553 sulfatase protein as shown in Figure 8.

30 Prosite program analysis was used to predict various sites within the 25278 sulfatase protein as shown in Figure 13.

Prosite program analysis was used to predict various sites within the 26212 sulfatase protein as shown in Figure 18.

In situ hybridization experiments showed that 22438 is expressed in subpopulations of DRG neurons, spinal cord, and brain, as disclosed hereinabove.

5 Expression of the 22438 sulfatase mRNA in the above cells and tissues indicates that the sulfatase is likely to be involved in the proper function of and in disorders involving these tissues. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment of sulfatase related disorders, especially disorders of these tissues that include, but are
10 not limited to those disclosed herein.

 The 23553 sulfatase is differentially expressed in breast and colon cancer and in colonic metastases to the liver. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment in these tissues (normal and tumor).

15 The 25278 sulfatase is differentially expressed in colon tumors and colonic metastases to the liver. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment in these normal and tumor tissues.

 The compositions include sulfatase polypeptides, nucleic acids, vectors,
20 transformed cells and related variants and fragments thereof, as well as agents that modulate expression of the polypeptides and polynucleotides. In particular, the invention relates to the modulation, diagnosis and treatment of sulfatase related disorders as described herein.

 Disorders involving the brain include, but are not limited to, disorders
25 involving neurons, and disorders involving glia, such as astrocytes, oligodendrocytes, ependymal cells, and microglia; cerebral edema, raised intracranial pressure and herniation, and hydrocephalus; malformations and developmental diseases, such as neural tube defects, forebrain anomalies, posterior fossa anomalies, and syringomyelia and hydromyelia; perinatal brain injury; cerebrovascular diseases, such as those
30 related to hypoxia, ischemia, and infarction, including hypotension, hypoperfusion, and low-flow states--global cerebral ischemia and focal cerebral ischemia--infarction from obstruction of local blood supply, intracranial hemorrhage, including

intracerebral (intraparenchymal) hemorrhage, subarachnoid hemorrhage and ruptured berry aneurysms, and vascular malformations, hypertensive cerebrovascular disease, including lacunar infarcts, slit hemorrhages, and hypertensive encephalopathy; infections, such as acute meningitis, including acute pyogenic (bacterial) meningitis and acute aseptic (viral) meningitis, acute focal suppurative infections, including brain abscess, subdural empyema, and extradural abscess, chronic bacterial meningoencephalitis, including tuberculosis and mycobacterioses, neurosyphilis, and neuroborreliosis (Lyme disease), viral meningoencephalitis, including arthropod-borne (Arbo) viral encephalitis, *Herpes simplex* virus Type 1, *Herpes simplex* virus Type 2, *Varicella-zoster* virus (*Herpes zoster*), cytomegalovirus, poliomyelitis, rabies, and human immunodeficiency virus 1, including HIV-1 meningoencephalitis (subacute encephalitis), vacuolar myelopathy, AIDS-associated myopathy, peripheral neuropathy, and AIDS in children, progressive multifocal leukoencephalopathy, subacute sclerosing panencephalitis, fungal meningoencephalitis, other infectious diseases of the nervous system; transmissible spongiform encephalopathies (prion diseases); demyelinating diseases, including multiple sclerosis, multiple sclerosis variants, acute disseminated encephalomyelitis and acute necrotizing hemorrhagic encephalomyelitis, and other diseases with demyelination; degenerative diseases, such as degenerative diseases affecting the cerebral cortex, including Alzheimer disease and Pick disease, degenerative diseases of basal ganglia and brain stem, including Parkinsonism, idiopathic Parkinson disease (paralysis agitans), progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy, including striatonigral degeneration, Shy-Drager syndrome, and olivopontocerebellar atrophy, and Huntington disease; spinocerebellar degenerations, including spinocerebellar ataxias, including Friedreich ataxia, and ataxia-telangiectasia, degenerative diseases affecting motor neurons, including amyotrophic lateral sclerosis (motor neuron disease), bulbospinal atrophy (Kennedy syndrome), and spinal muscular atrophy; inborn errors of metabolism, such as leukodystrophies, including Krabbe disease, metachromatic leukodystrophy, adrenoleukodystrophy, Pelizaeus-Merzbacher disease, and Canavan disease, mitochondrial encephalomyopathies, including Leigh disease and other mitochondrial encephalomyopathies; toxic and acquired metabolic diseases, including vitamin deficiencies such as thiamine (vitamin B₁) deficiency and

vitamin B₁₂ deficiency, neurologic sequelae of metabolic disturbances, including hypoglycemia, hyperglycemia, and hepatic encephatopathy, toxic disorders, including carbon monoxide, methanol, ethanol, and radiation, including combined methotrexate and radiation-induced injury; tumors, such as gliomas, including astrocytoma, including fibrillary (diffuse) astrocytoma and glioblastoma multiforme, pilocytic astrocytoma, pleomorphic xanthoastrocytoma, and brain stem glioma, oligodendroglioma, and ependymoma and related paraventricular mass lesions, neuronal tumors, poorly differentiated neoplasms, including medulloblastoma, other parenchymal tumors, including primary brain lymphoma, germ cell tumors, and pineal parenchymal tumors, meningiomas, metastatic tumors, paraneoplastic syndromes, peripheral nerve sheath tumors, including schwannoma, neurofibroma, and malignant peripheral nerve sheath tumor (malignant schwannoma), and neurocutaneous syndromes (phakomatoses), including neurofibromatosis, including Type 1 neurofibromatosis (NF1) and TYPE 2 neurofibromatosis (NF2), tuberous sclerosis, and Von Hippel-Lindau disease.

Furthermore, as disclosed in the background hereinabove, specific disorders have been associated with function of the various sulfatases. Accordingly, the sulfatases disclosed herein, having homology to specific sulfatases as disclosed herein, are useful for diagnosis and treatment of the disorders associated with sulfatase dysfunction as disclosed herein and to modulation of gene expression in the affected tissues.

The sequences of the invention find use in diagnosis of disorders involving an increase or decrease in sulfatase expression relative to normal expression, such as a proliferative disorder, a differentiative disorder, or a developmental disorder. The sequences also find use in modulating sulfatase-related responses. By "modulating" is intended the upregulating or downregulating of a response. That is, the compositions of the invention affect the targeted activity in either a positive or negative fashion.

The invention relates to novel sulfatases, having the deduced amino acid sequence shown in Figures 1, 5, 10, and 15 (SEQ ID NOS:1, 3, 5, and 7) or having the amino acid sequences encoded by the deposited cDNAs, ATCC Nos. _____. The deposited sequences, as well as the polypeptides encoded by the sequences, are

incorporated herein by reference and control in the event of any conflict, such as a sequencing error, with description in this application.

Thus, the present invention provides an isolated or purified sulfatase polypeptides and variants and fragments thereof. "Sulfatase polypeptide" or "sulfatase protein" refers to the polypeptide in SEQ ID NOS:1, 3, 5, or 7 or encoded by the deposited cDNAs. The term "sulfatase protein" or "sulfatase polypeptide," however, further includes the numerous variants described herein, as well as fragments derived from the full-length sulfatase and variants.

Sulfatase polypeptides can be purified to homogeneity. It is understood, however, that preparations in which the polypeptide is not purified to homogeneity are useful and considered to contain an isolated form of the polypeptide. The critical feature is that the preparation allows for the desired function of the polypeptide, even in the presence of considerable amounts of other components. Thus, the invention encompasses various degrees of purity.

As used herein, a polypeptide is said to be "isolated" or "purified" when it is substantially free of cellular material when it is isolated from recombinant and non-recombinant cells, or free of chemical precursors or other chemicals when it is chemically synthesized. A polypeptide, however, can be joined to another polypeptide with which it is not normally associated in a cell and still be considered "isolated" or "purified."

In one embodiment, the language "substantially free of cellular material" includes preparations of sulfatase having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the polypeptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20%, less than about 10%, or less than about 5% of the volume of the protein preparation.

The sulfatase polypeptide is also considered to be isolated when it is part of a membrane preparation or is purified and then reconstituted with membrane vesicles or liposomes.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the sulfatase polypeptide in which it is separated from chemical

precursors or other chemicals that are involved in its synthesis. The language
“substantially free of chemical precursors or other chemicals” includes, but is not limited
to, preparations of the polypeptide having less than about 30% (by dry weight) chemical
precursors or other chemicals, less than about 20% chemical precursors or other
5 chemicals, less than about 10% chemical precursors or other chemicals, or less than
about 5% chemical precursors or other chemicals.

In one embodiment, the sulfatase polypeptide comprises the amino acid sequence
shown in SEQ ID NOS:1, 3, 5, or 7. However, the invention also encompasses sequence
variants. Variants include a substantially homologous protein encoded by the same
10 genetic locus in an organism, i.e., an allelic variant. Variants also encompass proteins
derived from other genetic loci in an organism, but having substantial homology to the
sulfatase of SEQ ID NOS:1, 3, 5, or 7. Variants also include proteins substantially
homologous to the sulfatase but derived from another organism, i.e., an ortholog.
Variants also include proteins that are substantially homologous to the sulfatase that are
15 produced by chemical synthesis. Variants also include proteins that are substantially
homologous to the sulfatase that are produced by recombinant methods. It is
understood, however, that variants exclude any amino acid sequences disclosed prior to
the invention.

Preferred sulfatase polypeptides of the present invention have an amino acid
20 sequence sufficiently identical to the amino acid sequence of SEQ ID NOS:1, 3, 5, or
7. The term "sufficiently identical" is used herein to refer to a first amino acid or
nucleotide sequence that contains a sufficient or minimum number of identical or
equivalent (e.g., with a similar side chain) amino acid residues or nucleotides to a
second amino acid or nucleotide sequence such that the first and second amino acid or
25 nucleotide sequences have a common structural domain and/or common functional
activity. For example, amino acid or nucleotide sequences that contain a common
structural domain having at least about 45%, 50%, 55%, 60%, or 65% identity,
preferably 70-75% identity, more preferably 75-80%, 85-90%, 91%, 92%, 93%, 94%,
95%, 96%, 97%, 98% or 99% identity are defined herein as sufficiently identical.

30 In one embodiment, a variant of the 23553 sulfatase is greater than 92%
homologous. In another embodiment, a variant of the 25278 sulfatase is greater than

50% identical. In another embodiment, the 26212 sulfatase is greater than 50% identical.

The determination of percent identity between two sequences using the algorithms of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to the sulfatase nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to the sulfatase molecules of the invention. When utilizing BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

As used herein, two proteins (or a region of the proteins) are substantially homologous when the amino acid sequences are at least about 50-55%, 55-60%, 60-65%, 65-70%, 70-75%, 75-80%, typically at least about 80-85%, 85-90%, and most typically at least about 90-95% or more homologous. A substantially homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence hybridizing to the nucleic acid sequence, or portion thereof, of the sequence shown in SEQ ID NOS:2, 4, 6, or 8 under stringent conditions as more fully described below.

To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is

occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of
5 the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The invention also encompasses polypeptides having a lower degree of identity but having sufficient similarity so as to perform one or more of the same functions
10 performed by the sulfatase. Similarity is determined by conservative amino acid substitution, as shown in Table 1. Such substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Conservative substitutions are likely to be phenotypically silent. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala,
15 Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine
	Tryptophan
	Tyrosine
Hydrophobic	Leucine
	Isoleucine
	Valine
Polar	Glutamine
	Asparagine
Basic	Arginine
	Lysine
	Histidine
Acidic	Aspartic Acid
	Glutamic Acid
Small	Alanine
	Serine
	Threonine
	Methionine
	Glycine

A variant polypeptide can differ in amino acid sequence by one or more
 5 substitutions, deletions, insertions, inversions, fusions, and truncations or a
 combination of any of these. Variant polypeptides can be fully functional or can lack
 function in one or more activities. Thus, in the present case, variations can affect the
 function, for example, of one or more of regions including a metal (e.g., Ca^{++})-
 binding domain, activation domain, sulfatase catalytic domain, the region containing a
 10 propeptide, regulatory regions, substrate binding regions, regions involved in

membrane association or subcellular localization, regions involved in post-translational modification, for example, by phosphorylation, and regions that are important for effector function (i.e., agents that act upon the protein, such as in the conversion of cysteine to 2-amino-3-oxopropionic acid or serine semi-aldehyde).

5 Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Functional variants can also contain substitution of similar amino acids, which results in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

10 Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

As indicated, variants can be naturally-occurring or can be made by recombinant means or chemical synthesis to provide useful and novel characteristics for the sulfatase
15 polypeptide. This includes preventing immunogenicity from pharmaceutical formulations by preventing protein aggregation.

Useful variations further include alteration of functional activity. For example, one embodiment involves a variation at the substrate binding site that results in binding but not hydrolysis or more or less hydrolysis of the substrate than wild type. A further
20 useful variation at the same site can result in altered affinity for the substrate. Useful variations also include changes that provide for affinity for another substrate. Useful variations further include the ability to bind an effector molecule with greater or lesser affinity, such as not to bind or to bind but not release it. Further useful variations include alteration in the ability of the propeptide to be cleaved by a cleavage protein,
25 including alteration in the binding or recognition site. Further, the cleavage site can also be modified so that recognition and cleavage are by a different protease. A specific useful variation involves a variation in the ability to be bound or activated by the enzyme that activates the sulfatase by the conversion of cysteine to 2-3-oxopropionic acid or serine semi-aldehyde. Further variation could include a
30 variation in the specificity of metal binding.

Another useful variation provides a fusion protein in which one or more domains or subregions are operationally fused to one or more domains, subregions, or motifs

from another sulfatase. For example, a transmembrane domain from a protein can be introduced into the sulfatase such that the protein is anchored in the cell surface.

Other permutations include changing the number of sulfatase domains, and mixing of sulfatase domains from different sulfatase families, so that substrate specificity is altered. Mixing these various domains can allow the formation of novel sulfatase molecules with different host cell, subcellular localization, substrate, and effector molecule (one that acts on the sulfatase) specificity.

The term "substrate" is intended to refer not only to the sulfated substrate that is cleaved by the sulfatase domain, but to refer to any component with which the polypeptide interacts in order to produce an effect on that component or a subsequent biological effect that is a result of interacting with that component. This can include, but is not limited to, for example, interaction with the sulfatase activation enzyme and components involved in the conversion of 3' phosphoadenosine 5' phosphosulfate to adenosine 3' 5' biphosphate.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.* (1985) *Science* 244:1081-1085). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity, such as peptide bond hydrolysis *in vitro* or related biological activity, such as proliferative activity. Sites that are critical for binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.* (1992) *J. Mol. Biol.* 224:899-904; de Vos *et al.* (1992) *Science* 255:306-312).

The invention thus also includes polypeptide fragments of the sulfatases. Fragments can be derived from the amino acid sequence shown in SEQ ID NOS:1, 3, 5, or 7. However, the invention also encompasses fragments of the variants of the sulfatase polypeptides as described herein. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed prior to the present invention.

A fragment can comprise at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more contiguous amino acids. Fragments can retain one or more of the biological activities of the protein, for example as discussed

above, as well as fragments that can be used as an immunogen to generate sulfatase antibodies.

For example, for the 25278 sulfatase, the invention encompasses amino acid fragments greater than 5 amino acids, particularly from regions up to around
5 nucleotide 450 and beyond around nucleotide 1520. Specific fragments which may be excluded include those that are underlined in Figure 1. However, even in regions between around nucleotide 450 to around nucleotide 1520, fragments include those that are five or greater excluding those which may have been disclosed prior to the present invention.

10 For the 23553 sulfatase, fragments particularly include fragments of 5 amino acids or more up to around nucleotide 670.

For the 26212 sulfatase, for example, fragments containing 5 or more amino acids up to about nucleotide 572 are particularly encompassed by the invention. However, fragments of 5 amino acids or more encoded by around nucleotide 572 to
15 around nucleotide 1985 are also encompassed by the invention with the understanding that such fragments do not encompass those which may have been disclosed prior to the invention. For example, these can include the sections underlined in Figure 15.

Biologically active fragments (peptides which are, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 50, 100 or more amino acids in length) can comprise a
20 functional site. Such sites include but are not limited to those discussed above, such as a catalytic site, regulatory site, site important for substrate recognition or binding, regions containing a sulfatase domain or motif, phosphorylation sites, glycosylation sites, and other functional sites disclosed herein.

Fragments, for example, can extend in one or both directions from the functional
25 site to encompass 5, 10, 15, 20, 30, 40, 50, or up to 100 amino acids. Further, fragments can include sub-fragments of the specific sites or regions disclosed herein, which sub-fragments retain the function of the site or region from which they are derived.

The invention also provides fragments with immunogenic properties. These contain an epitope-bearing portion of the sulfatase polypeptide and variants. These
30 epitope-bearing peptides are useful to raise antibodies that bind specifically to a sulfatase polypeptide or region or fragment. These peptides can contain at least 10, 12, at least 14, or between at least about 15 to about 30 amino acids. The epitope-bearing sulfatase

polypeptides may be produced by any conventional means (Houghten, R.A. (1985) *Proc. Natl. Acad. Sci. USA* 82:5131-5135). Simultaneous multiple peptide synthesis is described in U.S. Patent No. 4,631,211.

5 Non-limiting examples of antigenic polypeptides that can be used to generate antibodies include but are not limited to peptides derived from extracellular regions. Regions having a high antigenicity index are shown in Figures 3, 7, 12, and 17. However, intracellularly-made antibodies ("intrabodies") are also encompassed, which would recognize intracellular peptide regions.

10 Fragments can be discrete (not fused to other amino acids or polypeptides) or can be within a larger polypeptide. Further, several fragments can be comprised within a single larger polypeptide. In one embodiment a fragment designed for expression in a host can have heterologous pre- and pro-polypeptide regions fused to the amino terminus of the sulfatase polypeptide fragment and an additional region fused to the carboxyl terminus of the fragment.

15 The invention thus provides chimeric or fusion proteins. These comprise a sulfatase peptide sequence operatively linked to a heterologous peptide having an amino acid sequence not substantially homologous to the sulfatase polypeptide. "Operatively linked" indicates that the sulfatase polypeptide and the heterologous peptide are fused in-frame. The heterologous peptide can be fused to the N-terminus or C-terminus of the
20 sulfatase polypeptide or can be internally located.

In one embodiment the fusion protein does not affect sulfatase function *per se*. For example, the fusion protein can be a GST-fusion protein in which sulfatase sequences are fused to the N- or C-terminus of the GST sequences. Other types of fusion proteins include, but are not limited to, enzymatic fusion proteins, for example
25 beta-galactosidase fusions, yeast two-hybrid GAL4 fusions, poly-His fusions and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant sulfatase polypeptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence. Therefore, in another embodiment, the fusion protein
30 contains a heterologous signal sequence at its C- or N-terminus.

EP-A-O 464 533 discloses fusion proteins comprising various portions of immunoglobulin constant regions. The Fc is useful in therapy and diagnosis and thus

results, for example, in improved pharmacokinetic properties (EP-A 0232 262). In drug discovery, for example, human proteins have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists (Bennett *et al.* (1995) *J. Mol. Recog.* 8:52-58 (1995) and Johanson *et al.* *J. Biol. Chem.* 270:9459-9471). Thus, this invention also encompasses soluble fusion proteins containing a sulfatase polypeptide and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclass (IgG, IgM, IgA, IgE). Preferred as immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. For some uses it is desirable to remove the Fc after the fusion protein has been used for its intended purpose, for example when the fusion protein is to be used as antigen for immunizations. In a particular embodiment, the Fc part can be removed in a simple way by a cleavage sequence, which is also incorporated and can be cleaved with factor Xa.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.* (1992) *Current Protocols in Molecular Biology*). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A sulfatase-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to sulfatase.

Another form of fusion protein is one that directly affects sulfatase functions. Accordingly, a sulfatase polypeptide is encompassed by the present invention in which one or more of the sulfatase regions (or parts thereof) has been replaced by heterologous or homologous regions (or parts thereof) from another sulfatase. Accordingly, various permutations are possible, for example, as discussed above. Thus, chimeric sulfatases can be formed in which one or more of the native domains or subregions has been

5 duplicated, removed, or replaced by another. This includes but is not limited to catalytic sulfatase or substrate binding domains, and regions involved in activation.

It is understood however that such regions could be derived from a sulfatase that has not yet been characterized. Moreover, sulfatase function can be derived from

5 peptides that contain these functions but are not in a sulfatase family.

The isolated 22438 sulfatase protein can be purified from cells that naturally express it, such as DRG neurons, including small and medium sized neurons, spinal cord, including interneurons and motor neurons, and brain, especially purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods.

10 The isolated 23553 sulfatase protein can be purified from cells that naturally express it, such as cells from any of the tissues shown in Figure 9, especially purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods.

15 The isolated 25278 sulfatase protein can be purified from cells that naturally express it, such as cells from any of the tissues shown in Figure 14, especially purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods.

The isolated 26212 sulfatase protein can be purified from cells that naturally express it, especially purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods.

20 In one embodiment, the protein is produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the sulfatase polypeptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques.

25 Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally-occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in

polypeptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art.

Accordingly, the polypeptides also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a
5 substituent group is included, in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence for purification of the mature polypeptide or a pro-protein sequence.

10 Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of
15 covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

20 Such modifications are well-known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd ed., T.E.
25 Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (1990) *Meth. Enzymol.* 182: 626-646 and Rattan *et al.* (1992) *Ann. N.Y. Acad. Sci.* 663:48-62).

30 As is also well known, polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of post-translation events, including

natural processing events and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translational natural processes and by synthetic methods.

Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. Blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally-occurring and synthetic polypeptides. For instance, the aminoterminal residue of polypeptides made in *E. coli*, prior to proteolytic processing, almost invariably will be N-formylmethionine.

The modifications can be a function of how the protein is made. For recombinant polypeptides, for example, the modifications will be determined by the host cell posttranslational modification capacity and the modification signals in the polypeptide amino acid sequence. Accordingly, when glycosylation is desired, a polypeptide should be expressed in a glycosylating host, generally a eukaryotic cell.

Insect cells often carry out the same posttranslational glycosylations as mammalian cells and, for this reason, insect cell expression systems have been developed to efficiently express mammalian proteins having native patterns of glycosylation. Similar considerations apply to other modifications.

The same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain more than one type of modification.

Polypeptide Uses

The protein sequences of the present invention can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped

alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

5 Sulfatase polypeptides are useful for producing antibodies specific for sulfatase, regions, or fragments. Regions having a high antigenicity index score are shown in Figures 3, 7, 12, and 17.

Sulfatase polypeptides are useful for biological assays related to sulfatases. Such assays involve any of the known sulfatase functions or activities or properties useful for
10 diagnosis and treatment of sulfatase-related conditions, including those in the references cited herein, which are incorporated by reference for these assays, functions, and disorders.

These assays include, but are not limited to, binding to and/or cleaving specific substrates to produce fragments, steady state levels of sulfated compounds, cysteine
15 modification, and biological assays related to the functions produced by sulfated compounds. Specific substrates useful for assays related to sulfate conjugate hydrolysis include but are not limited to xenobiotics, thyroid hormones, steroids, and catechols. Specific sulfate conjugates include, but are not limited to, 3 α -sulfatolithocholytaurine, sulfate conjugates of estrone, 4-methylumbelliferone, and harmol, sulfated cartilage and
20 proteoglycans, 4-nitrophenol, simple phenols, hydroxyarylamines, iodothyronines, catecholamines, 1-naphthyl, salbutamol, estrogens, ethinylestradiol, equilenin, diethylstilbestrol, androgens, cholesterol bile salts, pregnenolone, benzylic alcohols, glycolipidsulfates, complex carbohydrates such as dermatan and chondrotin sulfate, steroid sulfate, sulfate conjugates of xenobiotics, cholesterol sulfate, xenobiotic phenyls,
25 *o*-cresol, vanillin, eugenol, *m*-cresol, thymol, ethyl-4,4-dihydroxybenzoate, *p*-cresol, sesamol, methyl-2,6-dihydroxy-4-methylbenzyloate, methyl-2,4-dihydroxybenzoate, methyl-3,5-dihydroxybenzoate, tyramine, dopamine, 5 hydroxytryptamine, pyrogallol, 4-nitrocatecholsulfate, estrone sulfate, metabolites of the cytochrome P450 mono-oxygenase system, dihydroepiandrosterone sulfate (DHEAS), minoxidil, cicletanine,
30 sulfated mutagens and carcinogens, such as aromatic amines (including heterocyclic amines), and benzylic alcohols of chemicals such as polycyclic aromatic hydrocarbons, saffrole and estragole, glycosaminoglycans, sulfolipids, betahydroxysteroids, sulfate

esters of chromogenic or fluorogenic aromatic compounds, cerebroside sulfate, keratan sulfate, and heparan sulfate. Substrates also include any in the references cited herein, which are incorporated herein by reference for these substrates. Accordingly the assays include, but are not limited to, these sulfated substrates and biological effects of sulfation or desulfation of these substrates and associated biochemical, cellular, or phenotypic effects of sulfation of desulfation, and any of the other biological or functional properties of these proteins, including, but not limited to, those disclosed herein, and in any reference cited herein which is incorporated herein by reference for the disclosure of these properties and for the assays based on these properties. Further, assays may relate to changes in the protein, *per se*, and on the effects of these changes, for example, activation of the sulfatase by modification of a cysteine residue as disclosed herein, cleavage of the propeptide by a proteinase, induction of expression of the protein *in vivo*, inhibition of function, as well as any other effects on the protein mentioned herein or cited in any reference herein, which are incorporated herein by reference for these effects and for the subsequent biological consequences of these effects.

Sulfatase polypeptides are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express sulfatase, such as those discussed above, especially tumor cells, as a biopsy, or expanded in cell culture. In one embodiment, however, cell-based assays involve recombinant host cells expressing sulfatase. Accordingly, these drug-screening assays can be based on effects on protein function as described above for biological assays useful for diagnosis and treatment.

Determining the ability of the test compound to interact with a sulfatase can also comprise determining the ability of the test compound to preferentially bind to the polypeptide as compared to the ability of a known binding molecule to bind to the polypeptide.

The polypeptides can be used to identify compounds that modulate sulfatase activity. Such compounds, for example, can increase or decrease affinity or rate of binding to substrate, compete with substrate for binding to sulfatase, or displace substrate bound to sulfatase. Both sulfatase and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to sulfatase. These compounds can be further screened against a functional sulfatase to

determine the effect of the compound on sulfatase activity. Compounds can be identified that activate (agonist) or inactivate (antagonist) sulfatase to a desired degree. Modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject).

5 Sulfatase polypeptides can be used to screen a compound for the ability to stimulate or inhibit interaction between sulfatase protein and a target molecule that normally interacts with the sulfatase, for example, substrate of the sulfatase domain. The assay includes the steps of combining sulfatase protein with a candidate compound under conditions that allow the sulfatase protein or fragment to interact
10 with the target molecule, and to detect the formation of a complex between the sulfatase protein and the target or to detect the biochemical consequence of the interaction with the sulfatase and the target.

 Determining the ability of the sulfatase to bind to a target molecule can also be accomplished using a technology such as real-time Bimolecular Interaction Analysis
15 (BIA). Sjolander *et al.* (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological
20 molecules.

 The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-
25 compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

30 Examples of methods for the synthesis of molecular libraries can be found in the art, for example in DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med.*

Chem. 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233. Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on
5 beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 97:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-
10 310); (Ladner *supra*).

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.* (1991) *Nature* 354:82-84; Houghten *et al.* (1991) *Nature* 354:84-86) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration
15 amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.* (1993) *Cell* 72:767-778); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); 4) small organic and inorganic molecules (e.g.,
20 molecules obtained from combinatorial and natural product libraries); substrate analogs including, but not limited to, substrates disclosed herein.

One candidate compound is a soluble full-length sulfatase or fragment that competes for substrate. Other candidate compounds include mutant sulfatases or appropriate fragments containing mutations that affect sulfatase function and compete
25 for substrate. Accordingly, a fragment that competes for substrate, for example with a higher affinity, or a fragment that binds substrate but does not process or otherwise affect it, is encompassed by the invention.

The invention provides other end points to identify compounds that modulate (stimulate or inhibit) sulfatase activity. The assays typically involve an assay of cellular
30 events that indicate sulfatase activity. Thus, the expression of genes that are up- or down-regulated in response to sulfatase activity can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily

detectable, such as luciferase. Alternatively, modification of the sulfatase could also be measured.

Any of the biological or biochemical functions mediated by the sulfatase can be used as an endpoint assay. These include any of the biochemical or
5 biochemical/biological events described herein, in any reference cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art. Specific end points can include, but are not limited to, the events resulting from expression (or lack thereof) of sulfatase activity. With respect to disorders, this would include, but not be limited to, effects on function,
10 differentiation, and proliferation, which can be assayed, as well as the biological effects of function, such as disorders discussed hereinabove and in the references cited hereinabove which are incorporated herein by reference for the disorders disclosed in those references and other disorders and pathology. In the case of the 22438 sulfatase, models of pain can be used as an end point. In the case of the 23553
15 and 25278 sulfatases, tumor progression can be used as an end point.

Binding and/or activating compounds can also be screened by using chimeric sulfatase proteins in which one or more regions, segments, sites, and the like, as disclosed herein, or parts thereof, can be replaced by heterologous and homologous counterparts derived from other sulfatases. For example, a catalytic region can be used
20 that interacts with a different substrate specificity and/or affinity than the native sulfatase. Accordingly, a different set of components is available as an end-point assay for activation. As a further alternative, the site of modification by an effector protein, for example, activation or phosphorylation, can be replaced with the site for a different effector protein. Activation can also be detected by a reporter gene containing an easily
25 detectable coding region operably linked to a transcriptional regulatory sequence that is part of the native pathway in which sulfatase is involved.

Sulfatase polypeptides are also useful in competition binding assays in methods designed to discover compounds that interact with the sulfatase. Thus, a compound is exposed to a sulfatase polypeptide under conditions that allow the compound to bind or
30 to otherwise interact with the polypeptide. Soluble sulfatase polypeptide is also added to the mixture. If the test compound interacts with the soluble sulfatase polypeptide, it decreases the amount of complex formed or activity from the sulfatase target. This type

of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the sulfatase. Thus, the soluble polypeptide that competes with the target sulfatase region is designed to contain peptide sequences corresponding to the region of interest.

- 5 Another type of competition-binding assay can be used to discover compounds that interact with specific functional sites. As an example, bindable substrate analog and a candidate compound can be added to a sample of the sulfatase. Compounds that interact with the sulfatase at the same site as the substrate or analog will reduce the amount of complex formed between the sulfatase and the substrate or analog.
- 10 Accordingly, it is possible to discover a compound that specifically prevents interaction between the sulfatase and the component. Another example involves adding a candidate compound to a sample of sulfatase and cleavable substrate. A compound that competes with the substrate will reduce the amount of hydrolysis or binding of the substrate to the sulfatase. Accordingly, compounds can be discovered that directly interact with the
- 15 sulfatase and compete with the substrate. Such assays can involve any other component that interacts with the sulfatase.

To perform cell free drug screening assays, it is desirable to immobilize either sulfatase, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate

20 automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase/sulfatase fusion proteins can be adsorbed onto glutathione sepharose beads

25 (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ³⁵S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined

30 directly, or in the supernatant after the complexes is dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of sulfatase-binding protein found in the bead fraction quantitated from the gel using

standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be

5 derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a sulfatase-binding target component, such as substrate or activating enzyme, and a candidate compound are incubated in sulfatase-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized

10 complexes, include immunodetection of complexes using antibodies reactive with the sulfatase target molecule, or which are reactive with the sulfatase and compete with the target molecule; as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Modulators of sulfatase activity identified according to these drug screening

15 assays can be used to treat a subject with a disorder related to the sulfatase, by treating cells that express the sulfatase. These methods of treatment include the steps of administering the modulators of sulfatase activity in a pharmaceutical composition as described herein, to a subject in need of such treatment.

The 23553 and 25278 sulfatases are differentially expressed in tumor cells as

20 disclosed herein. Accordingly, these sulfatases are relevant to these disorders and relevant as well to differentiation, function, and growth of the tissues giving rise to the tumors. The 22438 sulfatase is expressed as described above, and accordingly is relevant for disorders involving these tissues. Disorders include, but are not limited to, those discussed hereinabove. Moreover, since the gene is expressed in the central

25 nervous system, this sulfatase is relevant for the treatment of pain.

Sulfatase polypeptides are thus useful for treating a sulfatase-associated disorder characterized by aberrant expression or activity of a sulfatase. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or

30 downregulates) expression or activity of the protein. In another embodiment, the method involves administering sulfatase as therapy to compensate for reduced or aberrant expression or activity of the protein.

Methods for treatment include but are not limited to the use of soluble sulfatase or fragments of sulfatase protein that compete for substrate or any other component that directly interacts with sulfatase, or any of the enzymes that modify the sulfatase. These sulfatases or fragments can have a higher affinity for the target so as to provide effective competition.

Stimulation of activity is desirable in situations in which the protein is abnormally downregulated and/or in which increased activity is likely to have a beneficial effect. Likewise, inhibition of activity is desirable in situations in which the protein is abnormally upregulated and/or in which decreased activity is likely to have a beneficial effect. In one example of such a situation, a subject has a disorder characterized by aberrant development or cellular differentiation. In another example, the subject has a disorder characterized by an aberrant hematopoietic response. In another example, it is desirable to achieve tissue regeneration in a subject.

In yet another aspect of the invention, the proteins of the invention can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO 94/10300), to identify other proteins (captured proteins) which bind to or interact with the proteins of the invention and modulate their activity.

Sulfatase polypeptides also are useful to provide a target for diagnosing a disease or predisposition to disease mediated by the sulfatase, including, but not limited to, those diseases disclosed herein, in the references cited herein, and as disclosed above in the background. Accordingly, methods are provided for detecting the presence, or levels of the sulfatase in a cell, tissue, or organism. The method involves contacting a biological sample with a compound capable of interacting with the sulfatase such that the interaction can be detected. One agent for detecting a sulfatase is an antibody capable of selectively binding to the sulfatase. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The sulfatase also provides a target for diagnosing active disease, or predisposition to disease, in a patient having a variant sulfatase. Thus, sulfatase can be

isolated from a biological sample and assayed for the presence of a genetic mutation that results in an aberrant protein. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered sulfatase activity in cell-based or cell-free assays, such as by alteration in substrate binding or degradation, or ability to be activated by the activation enzyme, or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein in general or in a sulfatase specifically, such as are disclosed herein.

In vitro techniques for detection of sulfatase include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. Alternatively, the protein can be detected *in vivo* in a subject by introducing into the subject a labeled anti-sulfatase antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods, which detect the allelic variant of sulfatase expressed in a subject, and methods, which detect fragments of sulfatase in a sample.

Sulfatase polypeptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985, and Linder, M.W. (1997) *Clin. Chem.* 43(2):254-266. The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes affects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the

expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of sulfatase in which one or more of sulfatase functions in one population is different from those in another population. The polypeptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a peptide-based treatment, polymorphism may give rise to catalytic regions that are more or less active. Accordingly, dosage would necessarily be modified to maximize the therapeutic effect within a given population containing the polymorphism. As an alternative to genotyping, specific polymorphic polypeptides could be identified.

Sulfatase polypeptides are also useful for monitoring therapeutic effects during clinical trials and other treatment. Thus, the therapeutic effectiveness of an agent that is designed to increase or decrease gene expression, protein levels or sulfatase activity can be monitored over the course of treatment using sulfatase polypeptides as an end-point target. The monitoring can be, for example, as follows: (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression or activity of the protein in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the protein in the post-administration samples; (v) comparing the level of expression or activity of the protein in the pre-administration sample with the protein in the post-administration sample or samples; and (vi) increasing or decreasing the administration of the agent to the subject accordingly.

Antibodies

The invention also provides antibodies that selectively bind to the sulfatase and its variants and fragments. An antibody is considered to selectively bind, even if it also binds to other proteins that are not substantially homologous with the sulfatase. These other proteins share homology with a fragment or domain of sulfatase. This conservation in specific regions gives rise to antibodies that bind to both proteins by

virtue of the homologous sequence. In this case, it would be understood that antibody binding to the sulfatase is still selective.

Antibodies can be polyclonal or monoclonal. An intact antibody, or a fragment thereof (e.g. Fab or F(ab')₂) can be used. An appropriate immunogenic preparation can
5 be derived from native, recombinantly expressed, or chemically synthesized peptides.

To generate antibodies, an isolated sulfatase polypeptide is used as an immunogen to generate antibodies using standard techniques for polyclonal and monoclonal antibody preparation. Either the full-length protein or antigenic peptide fragment can be used. Regions having a high antigenicity index are disclosed
10 hereinabove.

Antibodies are preferably prepared from these regions or from discrete fragments in these regions. However, antibodies can be prepared from any region of the peptide as described herein. A preferred fragment produces an antibody that diminishes or completely prevents substrate hydrolysis or binding. Antibodies can be
15 developed against the entire sulfatase or domains of the sulfatase as described herein, for example, the substrate binding region, sulfatase motif, or subregions thereof. Antibodies can also be developed against other specific functional sites as disclosed herein.

The antigenic peptide can comprise a contiguous sequence of at least 12, 14, 15-
20 20, 20-25, or 25-30 or more amino acid residues. In one embodiment, fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions. These fragments are not to be construed, however, as encompassing any fragments, which may be disclosed prior to the invention.

Detection can be facilitated by coupling (i.e., physically linking) the antibody to
25 a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin;
30 examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol;

examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Antibody Uses

5 The antibodies can be used to isolate a sulfatase by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural sulfatase from cells and recombinantly produced sulfatase expressed in host cells.

10 The antibodies are useful to detect the presence of a sulfatase in cells or tissues to determine the pattern of expression of the sulfatase among various tissues in an organism and over the course of normal development. The antibodies can be used to detect a sulfatase *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Antibody detection of circulating fragments of the full length sulfatase can be used to identify sulfatase turnover. In addition, the antibodies
15 can be used to assess abnormal tissue distribution or abnormal expression during development.

 Further, the antibodies can be used to assess sulfatase expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to sulfatase function. When a disorder is caused by an inappropriate
20 tissue distribution, developmental expression, or level of expression of sulfatase protein, the antibody can be prepared against the normal sulfatase protein. If a disorder is characterized by a specific mutation in sulfatase, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant sulfatase. However, intracellularly-made antibodies ("intrabodies") are also encompassed, which would
25 recognize intracellular sulfatase peptide regions.

 The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Antibodies can be developed against the whole sulfatase or portions of the sulfatase.

30 The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting sulfatase expression level or the presence of aberrant sulfatases and aberrant

tissue distribution or developmental expression, antibodies directed against the sulfatase or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic sulfatase can be used to identify individuals
5 that require modified treatment modalities.

The antibodies are also useful as diagnostic tools as an immunological marker for aberrant sulfatase analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Thus, where a specific sulfatase
10 has been correlated with expression in a specific tissue, antibodies that are specific for this sulfatase can be used to identify a tissue type.

The antibodies are also useful in forensic identification. Accordingly, where an individual has been correlated with a specific genetic polymorphism resulting in a specific polymorphic protein, an antibody specific for the polymorphic protein can be
15 used as an aid in identification.

The antibodies are also useful for inhibiting sulfatase function, for example, substrate binding, or sulfatase activity. For example, sulfatase activity may be measured by the ability to form a binding complex with a sulfated conjugate, such as disclosed herein.

20 These uses can also be applied in a therapeutic context in which treatment involves inhibiting sulfatase function. An antibody can be used, for example, to block substrate binding. Antibodies can be prepared against specific fragments containing sites required for function or against intact sulfatase associated with a cell.

Completely human antibodies are particularly desirable for therapeutic treatment
25 of human patients. For an overview of this technology for producing human antibodies, see Lonberg *et al.* (1995) *Int. Rev. Immunol.* 13:65-93. For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, e.g., U.S. Patent 5,625,126; U.S. Patent 5,633,425; U.S. Patent 5,569,825; U.S. Patent 5,661,016; and U.S. Patent
30 5,545,806.

The invention also encompasses kits for using antibodies to detect the presence of a sulfatase protein in a biological sample. The kit can comprise antibodies such as a

labeled or labelable antibody and a compound or agent for detecting the sulfatase in a biological sample; means for determining the amount of sulfatase in the sample; and means for comparing the amount of sulfatase in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further
5 comprise instructions for using the kit to detect the sulfatase.

Polynucleotides

The nucleotide sequences in SEQ ID NOS:2, 4, 6, and 8 were obtained by sequencing the deposited human cDNAs. Accordingly, the sequences of the deposited
10 clones are controlling as to any discrepancies between the two and any reference to a sequence of SEQ ID NOS:2, 4, 6, or 8, includes reference to the sequence of the deposited cDNA.

The specifically disclosed cDNA comprises the coding region and 5' and 3' untranslated sequences in SEQ ID NOS:2, 4, 6, or 8.

15 The invention provides isolated polynucleotides encoding the novel sulfatases. The term "sulfatase polynucleotide" or "sulfatase nucleic acid" refers to the sequences shown in SEQ ID NOS:2, 4, 6, or 8, or in the deposited cDNAs. The term "sulfatase polynucleotide" or "sulfatase nucleic acid" further includes variants and fragments of sulfatase polynucleotides.

20 An "isolated" sulfatase nucleic acid is one that is separated from other nucleic acid present in the natural source of sulfatase nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank sulfatase nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking
25 nucleotide sequences, for example up to about 5KB. The important point is that the sulfatase nucleic acid is isolated from flanking sequences such that it can be subjected to the specific manipulations described herein, such as recombinant expression, preparation of probes and primers, and other uses specific to the sulfatase nucleic acid sequences. In one embodiment, the sulfatase nucleic acid comprises only the coding region.

30 Moreover, an "isolated" nucleic acid molecule, such as a cDNA or RNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when

chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

In some instances, the isolated material will form part of a composition (for example, a crude extract containing other substances), buffer system or reagent mix.

5 In other circumstances, the material may be purified to essential homogeneity, for example as determined by PAGE or column chromatography such as HPLC.

Preferably, an isolated nucleic acid comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present.

For example, recombinant DNA molecules contained in a vector are considered
10 isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced
15 synthetically.

In some instances, the isolated material will form part of a composition (or example, a crude extract containing other substances), buffer system or reagent mix. In other circumstances, the material may be purified to essential homogeneity, for example as determined by PAGE or column chromatography such as HPLC. Preferably, an
20 isolated nucleic acid comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present.

Sulfatase polynucleotides can encode the mature protein plus additional amino or carboxyterminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences
25 may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

30 Sulfatase polynucleotides include, but are not limited to, the sequence encoding the mature polypeptide alone, the sequence encoding the mature polypeptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or

pro-protein sequence), the sequence encoding the mature polypeptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the polynucleotide may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Sulfatase polynucleotides can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides variant sulfatase polynucleotides, and fragments thereof, that differ from the nucleotide sequence shown in SEQ ID NOS:2, 4, 6, or 8, due to degeneracy of the genetic code and thus encode the same protein as that encoded by a nucleotide sequence shown in SEQ ID NOS:2, 4, 6, or 8.

The invention also provides sulfatase nucleic acid molecules encoding the variant polypeptides described herein. Such polynucleotides may be naturally occurring, such as allelic variants (same locus), homologs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions.

Typically, variants have a substantial identity with a nucleic acid molecules of SEQ ID NOS:2, 4, 6 or 8, and the complements thereof. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. These variants comprise a nucleotide sequence encoding a sulfatase that is typically at least about 40-45%, 45-50%, 50-55%, 55-60%, 60-65%, 65-70%, 70-75%, more typically at least about 75-80% or 80-85%, and most typically at least about

85-90% or 90-95% or more homologous to the nucleotide sequence shown in SEQ ID NOS:2, 4, 6 or 8, or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under stringent conditions, to the nucleotide sequence shown in SEQ ID NOS:2, 4, 6 or 8, or a fragment of the sequence.

5 In the case of the 23553 sulfatase, in one embodiment, a variant is greater than 65% homologous with respect to nucleotide sequence. For the 25278 sulfatase, in one embodiment, a variant is greater than 50-60% homologous with respect to nucleotide sequence. With respect to the 26212 sulfatase, in one embodiment, a variant is greater than about 65-75% homologous with respect to nucleotide sequence.

10 It is understood that stringent hybridization does not indicate substantial homology where it is due to general homology, such as polyA⁺ sequences, or sequences common to all or most proteins, sulfatases, arylsulfatases, glucosamine-6-sulfatases, N-acetylgalactosamine-4-sulfatases, or any of the sulfatases to which the sulfatases of the present invention have shown homology by BLAST analysis, for example, regions to
15 arylsulfatases A, B, C, D, E, F, IDS, and the like. Moreover, it is understood that variants do not include any of the nucleic acid sequences that may have been disclosed prior to the invention.

As used herein, the term "stringent conditions" is intended to describe conditions comprising hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C,
20 followed by one or more washes in 0.2X SSC/0.1% SDS at 65°C. Methods of hybridization are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1998), incorporated by reference. In one embodiment, an isolated nucleic acid molecule that hybridizes under stringent conditions to a sequence of SEQ ID NOS:2, 4, 6 or 8, corresponds to a naturally-
25 occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

The present invention also provides isolated nucleic acids that contain a single or double stranded fragment or portion that hybridizes under stringent conditions to
30 the nucleotide sequence of SEQ ID NOS:2, 4, 6, or 8 or the complements of SEQ ID NOS:2, 4, 6, or 8. In one embodiment, the nucleic acid consists of a portion of a nucleotide sequence of SEQ ID NOS:2, 4, 6 or 8, and the complements. The nucleic

acid fragments of the invention are at least about 10-15, preferably at least about 15-20 or 20-25 contiguous nucleotides, and can be 30, 33, 35, 40, 50, 60, 70, 75, 80, 90, 100, 200, 500 or more nucleotides in length. Longer fragments, for example, 600 or more nucleotides in length, which encode antigenic proteins or polypeptides described
5 herein are also useful.

In the case of the 23553 sulfatase, in one embodiment, fragments are derived from nucleotide 1 to about nucleotide 670 and comprise 5-10 and 10-20 contiguous base pairs, and particularly greater than 18. For this sulfatase, in another embodiment, a fragment is derived from around nucleotide 3008 to 3514 and
10 comprises around 5-10 and 10-20 contiguous nucleotides. In other embodiments for this sulfatase, a fragment is derived from around nucleotide 3994 to 4321 and is about 5-10 or 10-20 contiguous nucleotides. For the 25278, in one embodiment, a fragment is derived from around nucleotide 130 to around nucleotide 454 and comprises a contiguous sequence of about 5-10 or 10-20 nucleotides. In another embodiment, the
15 fragment is derived from around nucleotide 454 to around nucleotide 1400 and comprises around 5-10 or 10-20 contiguous nucleotides, especially a fragment greater than 17 nucleotides. In another embodiment the fragment is derived from around nucleotide 1400 to around nucleotide 1850 and comprises a continuous sequence of around 5-10, 10-20, or 20-25 nucleotides, especially a fragment greater than 23
20 nucleotides. In another embodiment, a fragment is derived from about nucleotide 1933 to about nucleotide 2421. Such a fragment comprises around 5-10 or 10-20 contiguous nucleotides. For the 26212 sulfatase, in one embodiment, a fragment is derived from around nucleotide 272 to around nucleotide 538 and comprises a contiguous sequence of around 5-10 or 10-20 nucleotides, especially a fragment
25 greater than 17 nucleotides. In another embodiment, the fragment is derived from around nucleotide 538 to around nucleotide 751 and comprises a contiguous sequence of at least 5-10 or 10-20 nucleotides, especially greater than 12 nucleotides. In another embodiment, the fragment is derived from around nucleotide 1074 to around 1551 and comprises a contiguous nucleotide sequence of around 5-10, 10-20, or 20-
30 30, especially greater than 20 nucleotides. In a further embodiment, the fragment is derived from around nucleotide 2052 to 2251 and comprises a contiguous sequence of 5-10 and 10-20 nucleotides, especially fragments greater than 18 nucleotides.

The fragment can comprise DNA or RNA and can be derived from either the coding or the non-coding sequence.

In another embodiment an isolated sulfatase nucleic acid encodes the entire coding region. In another embodiment the isolated sulfatase nucleic acid encodes a sequence corresponding to the mature protein. Other fragments include nucleotide sequences encoding the amino acid fragments described herein.

Thus, sulfatase nucleic acid fragments further include sequences corresponding to the regions described herein, subregions also described, and specific functional sites. Sulfatase nucleic acid fragments also include combinations of the regions, segments, motifs, and other functional sites described above. It is understood that a sulfatase fragment includes any nucleic acid sequence that does not include the entire gene. A person of ordinary skill in the art would be aware of the many permutations that are possible. Nucleic acid fragments, according to the present invention, are not to be construed as encompassing those fragments that may have been disclosed prior to the invention.

Where the location of the regions or sites have been predicted by computer analysis, one of ordinary skill would appreciate that the amino acid residues constituting these regions can vary depending on the criteria used to define the regions.

Polynucleotide Uses

The nucleotide sequences of the present invention can be used as a "query sequence" to perform a search against public databases, for example, to identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The nucleic acid fragments of the invention provide probes or primers in assays such as those described below. "Probes" are oligonucleotides that hybridize in a base-specific manner to a complementary strand of nucleic acid. Such probes include polypeptide nucleic acids, as described in Nielsen *et al.* (1991) *Science* 254:1497-1500. Typically, a probe comprises a region of nucleotide sequence that hybridizes under highly stringent conditions to at least about 15, typically about 20-25, and more typically about 30, 40, 50 or 75 consecutive nucleotides of the nucleic acid sequence shown in SEQ ID NO:2 and the complements thereof. More typically, the probe further comprises a label, e.g., radioisotope, fluorescent compound, enzyme, or enzyme co-factor.

As used herein, the term "primer" refers to a single-stranded oligonucleotide which acts as a point of initiation of template-directed DNA synthesis using well-known methods (e.g., PCR, LCR) including, but not limited to those described herein. The appropriate length of the primer depends on the particular use, but typically ranges from about 15 to 30 nucleotides. The term "primer site" refers to the area of the target DNA to which a primer hybridizes. The term "primer pair" refers to a set of primers including a 5' (upstream) primer that hybridizes with the 5' end of the nucleic acid sequence to be amplified and a 3' (downstream) primer that hybridizes with the complement of the sequence to be amplified.

Sulfatase polynucleotides are thus useful for probes, primers, and in biological assays. Where the polynucleotides are used to assess sulfatase properties or functions, such as in the assays described herein, all or less than all of the entire cDNA can be useful. Assays specifically directed to sulfatase functions, such as assessing agonist or antagonist activity, encompass the use of known fragments. Further, diagnostic methods for assessing sulfatase function can also be practiced with any fragment, including those fragments that may have been known prior to the invention. Similarly, in methods involving treatment of sulfatase dysfunction, all fragments are encompassed including those, which may have been known in the art.

Sulfatase polynucleotides are useful as a hybridization probe for cDNA and genomic DNA to isolate a full-length cDNA and genomic clones encoding the polypeptides described in SEQ ID NOS:1, 3, 5, or 7, and to isolate cDNA and genomic clones that correspond to variants producing the same polypeptides shown in SEQ ID

NOS:1, 3, 5, or 7, or the other variants described herein. Variants can be isolated from the same tissue and organism from which a polypeptide shown in SEQ ID NOS:1, 3, 5, or 7 was isolated, different tissues from the same organism, or from different organisms. This method is useful for isolating genes and cDNA that are developmentally-controlled and therefore may be expressed in the same tissue or different tissues at different points in the development of an organism.

The probe can correspond to any sequence along the entire length of the gene encoding the sulfatase polypeptide. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions.

The nucleic acid probe can be, for example, the full-length cDNA of SEQ ID NOS:2, 4, 6, or 8, or a fragment thereof, such as an oligonucleotide of at least 5, 10, 15, 20, 25, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to mRNA or DNA.

Fragments of the polynucleotides described herein are also useful to synthesize larger fragments or full-length polynucleotides described herein, ribozymes or antisense molecules. For example, a fragment can be hybridized to any portion of an mRNA and a larger or full-length cDNA can be produced.

Antisense nucleic acids of the invention can be designed using the nucleotide sequences of SEQ ID NOS:2, 4, 6, or 8, and constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-

methyaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest).

Additionally, the nucleic acid molecules of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4:5). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, e.g., DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996), *supra*; Perry-O'Keefe *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:14670. PNAs can be further modified, e.g., to enhance their stability, specificity or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996), *supra*, Finn *et al.* (1996) *Nucleic Acids Res.* 24(17):3357-63, Mag *et al.* (1989) *Nucleic Acids Res.* 17:5973, and Peterser *et al.* (1975) *Bioorganic Med. Chem. Lett.* 5:1119.

The nucleic acid molecules and fragments of the invention can also include other appended groups such as peptides (e.g., for targeting host cell sulfatases *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger *et*

al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. WO 88/0918) or the blood brain barrier (see, e.g., PCT Publication No. WO 89/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (see, e.g., Krol *et al.* (1988) *Bio-Techniques* 6:958-976) or intercalating agents (see, e.g., Zon (1988) *Pharm Res.* 5:539-549).

Sulfatase polynucleotides are also useful as primers for PCR to amplify any given region of a sulfatase polynucleotide.

Sulfatase polynucleotides are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the sulfatase polypeptides. Vectors also include insertion vectors, used to integrate into another polynucleotide sequence, such as into the cellular genome, to alter *in situ* expression of sulfatase genes and gene products. For example, an endogenous sulfatase coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

Sulfatase polynucleotides are also useful for expressing antigenic portions of sulfatase proteins.

Sulfatase polynucleotides are also useful as probes for determining the chromosomal positions of sulfatase polynucleotides by means of *in situ* hybridization methods, such as FISH. (For a review of this technique, see Verma *et al.* (1988) *Human Chromosomes: A Manual of Basic Techniques* (Pergamon Press, New York), and PCR mapping of somatic cell hybrids. The mapping of the sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, *Mendelian Inheritance in*

Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland *et al.* ((1987) *Nature* 325:783-787).

5 Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with a specified gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for
10 structural alterations in the chromosomes, such as deletions or translocations, that are visible from chromosome spreads, or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

15 Sulfatase polynucleotide probes are also useful to determine patterns of the presence of the gene encoding sulfatases and their variants with respect to tissue distribution, for example, whether gene duplication has occurred and whether the duplication occurs in all or only a subset of tissues. The genes can be naturally occurring or can have been introduced into a cell, tissue, or organism exogenously.

20 Sulfatase polynucleotides are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from genes encoding the polynucleotides described herein.

 Sulfatase polynucleotides are also useful for constructing host cells expressing a part, or all, of a sulfatase polynucleotide or polypeptide.

25 Sulfatase polynucleotides are also useful for constructing transgenic animals expressing all, or a part, of a sulfatase polynucleotide or polypeptide.

 Sulfatase polynucleotides are also useful for making vectors that express part, or all, of a sulfatase polypeptide.

 Sulfatase polynucleotides are also useful as hybridization probes for determining
30 the level of sulfatase nucleic acid expression. Accordingly, the probes can be used to detect the presence of, or to determine levels of, sulfatase nucleic acid in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA.

Accordingly, probes corresponding to the polypeptides described herein can be used to assess gene copy number in a given cell, tissue, or organism. This is particularly relevant in cases in which there has been an amplification of a sulfatase gene.

Alternatively, the probe can be used in an *in situ* hybridization context to assess the position of extra copies of a sulfatase gene, as on extrachromosomal elements or as integrated into chromosomes in which the sulfatase gene is not normally found, for example, as a homogeneously staining region.

These uses are relevant for diagnosis of disorders involving an increase or decrease in sulfatase expression relative to normal, such as a proliferative disorder, a differentiative or developmental disorder, or a hematopoietic disorder. Disorders in which sulfatase expression is relevant include, but are not limited to, those disclosed herein above.

Disorders in which 22438 sulfatase expression is relevant include, but are not limited to, those involving the tissues as disclosed herein and those associated with pain.

Disorders in which 23553 sulfatase expression is relevant include, but are not limited to, breast and colon carcinoma.

Disorders in which 25278 sulfatase expression is relevant include, but are not limited to, colon carcinoma.

Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant expression or activity of a sulfatase nucleic acid, in which a test sample is obtained from a subject and nucleic acid (e.g., mRNA, genomic DNA) is detected, wherein the presence of the nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant expression or activity of the nucleic acid.

One aspect of the invention relates to diagnostic assays for determining nucleic acid expression as well as activity in the context of a biological sample (e.g., blood, serum, cells, tissue) to determine whether an individual has a disease or disorder, or is at risk of developing a disease or disorder, associated with aberrant nucleic acid expression or activity. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of

a disorder characterized by or associated with expression or activity of the nucleic acid molecules.

In vitro techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern

5 hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a sulfatase, such as by measuring the level of a sulfatase-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if the sulfatase gene has been mutated.

10 Nucleic acid expression assays are useful for drug screening to identify compounds that modulate sulfatase nucleic acid expression (e.g., antisense, polypeptides, peptidomimetics, small molecules or other drugs). A cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of the mRNA in the presence of the candidate compound is compared to the
15 level of expression of the mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. The modulator can bind to the nucleic acid or indirectly modulate expression, such as by interacting with other cellular components
20 that affect nucleic acid expression.

Modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject) in patients or in transgenic animals. The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of
25 a sulfatase gene. The method typically includes assaying the ability of the compound to modulate the expression of the sulfatase nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired sulfatase nucleic acid expression.

The assays can be performed in cell-based and cell-free systems. Cell-based
30 assays include cells naturally expressing the sulfatase nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences. Alternatively, candidate compounds can be assayed *in vivo* in patients or in transgenic animals.

The assay for sulfatase nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds (such as substrate hydrolysis). Further, the expression of genes that are up- or down-regulated in response to sulfatase activity can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of sulfatase gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of sulfatase mRNA in the presence of the candidate compound is compared to the level of expression of sulfatase mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

Accordingly, the invention provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate sulfatase nucleic acid expression. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or effects on nucleic acid activity (e.g. when nucleic acid is mutated or improperly modified). Treatment is of disorders characterized by aberrant expression or activity of the nucleic acid.

Alternatively, a modulator for sulfatase nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits sulfatase nucleic acid expression.

Sulfatase polynucleotides are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of a sulfatase gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can

also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a
5 desirable level, administration of the compound could be commensurately decreased.

Monitoring can be, for example, as follows: (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a specified mRNA or genomic DNA of the invention in the pre-administration sample; (iii) obtaining one or more post-administration samples from
10 the subject; (iv) detecting the level of expression or activity of the mRNA or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the mRNA or genomic DNA in the pre-administration sample with the mRNA or genomic DNA in the post-administration sample or samples; and (vi) increasing or decreasing the administration of the agent to the subject accordingly.

15 Sulfatase polynucleotides are also useful in diagnostic assays for qualitative changes in sulfatase nucleic acid, and particularly in qualitative changes that lead to pathology. The polynucleotides can be used to detect mutations in sulfatase genes and gene expression products such as mRNA. The polynucleotides can be used as hybridization probes to detect naturally-occurring genetic mutations in a sulfatase gene
20 and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of a
25 sulfatase gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a sulfatase.

Mutations in a sulfatase gene can be detected at the nucleic acid level by a variety of techniques. Genomic DNA can be analyzed directly or can be amplified by
30 using PCR prior to analysis. RNA or cDNA can be used in the same way.

In certain embodiments, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195

and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.* (1988) *Bio/Technology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well-known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

Alternatively, mutations in a sulfatase gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method.

Furthermore, sequence differences between a mutant sulfatase gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated
5 sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the gene include methods in which
10 protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.* (1985) *Science* 230:1242); Cotton *et al.* (1988) *PNAS* 85:4397; Saleeba *et al.* (1992) *Meth. Enzymol.* 217:286-295), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.* (1989) *PNAS* 86:2766; Cotton *et al.* (1993) *Mutat. Res.* 285:125-144; and Hayashi *et al.* (1992) *Genet. Anal.*
15 *Tech. Appl.* 9:73-79), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.* (1985) *Nature* 313:495). The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject
20 method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet.* 7:5). Examples of other techniques for detecting point mutations include, selective oligonucleotide hybridization, selective amplification, and selective primer extension.

25 In other embodiments, genetic mutations can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotide probes (Cronin *et al.* (1996) *Human Mutation* 7:244-255; Kozal *et al.* (1996) *Nature Medicine* 2:753-759). For example, genetic mutations can be identified in two dimensional arrays containing
30 light-generated DNA probes as described in Cronin *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear

arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is
5 composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

Sulfatase polynucleotides are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the polynucleotides can be used to study the relationship between an
10 individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). In the present case, for example, a mutation in the sulfatase gene that results in altered affinity for a substrate-related compound could result in an excessive or decreased drug effect with standard concentrations of the compound. Accordingly, the sulfatase polynucleotides described herein can be used to
15 assess the mutation content of the gene in an individual in order to select an appropriate compound or dosage regimen for treatment.

Thus polynucleotides displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow
20 effective clinical design of treatment compounds and dosage regimens.

The methods can involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting mRNA, or genomic DNA, such that the presence of mRNA or genomic DNA is detected in the biological sample, and comparing the presence of mRNA or genomic
25 DNA in the control sample with the presence of mRNA or genomic DNA in the test sample.

Sulfatase polynucleotides are also useful for chromosome identification when the sequence is identified with an individual chromosome and to a particular location on the chromosome. First, the DNA sequence is matched to the chromosome by *in situ* or
30 other chromosome-specific hybridization. Sequences can also be correlated to specific chromosomes by preparing PCR primers that can be used for PCR screening of somatic cell hybrids containing individual chromosomes from the desired species. Only hybrids

containing the chromosome containing the gene homologous to the primer will yield an amplified fragment. Sublocalization can be achieved using chromosomal fragments. Other strategies include prescreening with labeled flow-sorted chromosomes and preselection by hybridization to chromosome-specific libraries. Further mapping strategies include fluorescence *in situ* hybridization, which allows hybridization with probes shorter than those traditionally used. Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on the chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Sulfatase polynucleotides can also be used to identify individuals from small biological samples. This can be done for example using restriction fragment-length polymorphism (RFLP) to identify an individual. Thus, the polynucleotides described herein are useful as DNA markers for RFLP (See U.S. Patent No. 5,272,057).

Furthermore, the sulfatase sequences can be used to provide an alternative technique, which determines the actual DNA sequence of selected fragments in the genome of an individual. Thus, the sulfatase sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify DNA from an individual for subsequent sequencing.

Panels of corresponding DNA sequences from individuals prepared in this manner can provide unique individual identifications, as each individual will have a unique set of such DNA sequences. It is estimated that allelic variation in humans occurs with a frequency of about once per each 500 bases. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. Sulfatase sequences can be used to obtain such identification sequences from individuals and from tissue. The sequences represent unique fragments of the human genome. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes.

If a panel of reagents from the sequences is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

Sulfatase polynucleotides can also be used in forensic identification procedures. PCR technology can be used to amplify DNA sequences taken from very small biological samples, such as a single hair follicle, body fluids (e.g. blood, saliva, or semen). The amplified sequence can then be compared to a standard allowing identification of the origin of the sample.

Sulfatase polynucleotides can thus be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e. another DNA sequence that is unique to a particular individual). As described above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to the noncoding region are particularly useful since greater polymorphism occurs in the noncoding regions, making it easier to differentiate individuals using this technique.

Sulfatase polynucleotides can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue. This is useful in cases in which a forensic pathologist is presented with a tissue of unknown origin. Panels of sulfatase probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these primers and probes can be used to screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

Alternatively, sulfatase polynucleotides can be used directly to block transcription or translation of sulfatase gene sequences by means of antisense or ribozyme constructs. Thus, in a disorder characterized by abnormally high or undesirable sulfatase gene expression, nucleic acids can be directly used for treatment.

Sulfatase polynucleotides are thus useful as antisense constructs to control sulfatase gene expression in cells, tissues, and organisms. A DNA antisense polynucleotide is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of sulfatase protein. An antisense RNA or DNA polynucleotide would hybridize to the mRNA and thus block translation of mRNA into sulfatase protein.

Examples of antisense molecules useful to inhibit nucleic acid expression include antisense molecules complementary to a fragment of the 5' untranslated region of SEQ ID NOS:2, 4, 6, or 8, which also includes the start codon and antisense molecules which are complementary to a fragment of the 3' untranslated region of SEQ ID NOS:2, 4, 6, or 8.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of sulfatase nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired sulfatase nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the sulfatase protein.

Sulfatase polynucleotides also provide vectors for gene therapy in patients containing cells that are aberrant in sulfatase gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired sulfatase protein to treat the individual.

The invention also encompasses kits for detecting the presence of a sulfatase nucleic acid in a biological sample. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting sulfatase nucleic acid in a biological sample; means for determining the amount of sulfatase nucleic acid in the sample; and means for comparing the amount of sulfatase nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect sulfatase mRNA or DNA.

Computer Readable Means

The nucleotide or amino acid sequences of the invention are also provided in a variety of mediums to facilitate use thereof. As used herein, "provided" refers to a manufacture, other than an isolated nucleic acid or amino acid molecule, which
5 contains a nucleotide or amino acid sequence of the present invention. Such a manufacture provides the nucleotide or amino acid sequences, or a subset thereof (e.g., a subset of open reading frames (ORFs)) in a form which allows a skilled artisan to examine the manufacture using means not directly applicable to examining the nucleotide or amino acid sequences, or a subset thereof, as they exists in nature or in
10 purified form.

In one application of this embodiment, a nucleotide or amino acid sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage
15 media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. The skilled artisan will readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium
20 having recorded thereon a nucleotide or amino acid sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable medium. The skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising the nucleotide or amino acid sequence information of the
25 present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide or amino acid sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition,
30 a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in

commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. The skilled artisan can readily adapt any number of dataprocessor structuring formats (e.g., text file or database) in order to obtain
5 computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing the nucleotide or amino acid sequences of the invention in computer readable form, the skilled artisan can routinely access the sequence information for a variety of purposes. For example, one skilled in the art can use the
10 nucleotide or amino acid sequences of the invention in computer readable form to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif.

As used herein, a "target sequence" can be any DNA or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300
15 nucleotide residues. However, it is well recognized that commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are
25 chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

Computer software is publicly available which allows a skilled artisan to
30 access sequence information provided in a computer readable medium for analysis and comparison to other sequences. A variety of known algorithms are disclosed

publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, MacPattern (EMBL), BLASTN and BLASTX (NCBIA).

5 For example, software which implements the BLAST (Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410) and BLAZE (Brutlag *et al.* (1993) *Comp. Chem.* 17:203-207) search algorithms on a Sybase system can be used to identify open reading frames (ORFs) of the sequences of the invention which contain homology to ORFs or proteins from other libraries. Such ORFs are protein encoding fragments and are
10 useful in producing commercially important proteins such as enzymes used in various reactions and in the production of commercially useful metabolites.

Vectors/Host Cells

 The invention also provides vectors containing sulfatase polynucleotides. The
15 term "vector" refers to a vehicle, preferably a nucleic acid molecule that can transport sulfatase polynucleotides. When the vector is a nucleic acid molecule, the sulfatase polynucleotides are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC,
20 PAC, YAC, OR MAC.

 A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of sulfatase polynucleotides. Alternatively, the vector may integrate into the host cell genome and produce additional copies of sulfatase polynucleotides when the host cell replicates.

25 The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of sulfatase polynucleotides. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

 Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to sulfatase polynucleotides such that transcription of the polynucleotides is
30 allowed in a host cell. The polynucleotides can be introduced into the host cell with a separate polynucleotide capable of affecting transcription. Thus, the second polynucleotide may provide a trans-acting factor interacting with the cis-regulatory

control region to allow transcription of sulfatase polynucleotides from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself.

5 It is understood, however, that in some embodiments, transcription and/or translation of sulfatase polynucleotides can occur in a cell-free system.

The regulatory sequence to which the polynucleotides described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate
10 early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early
15 enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as
20 polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual 2nd. ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

25 A variety of expression vectors can be used to express a sulfatase polynucleotide. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses,
30 poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression

vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual 2nd. ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

5 The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

10 Sulfatase polynucleotides can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

15 The vector containing the appropriate polynucleotide can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

20 As described herein, it may be desirable to express the polypeptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of sulfatase polypeptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic
25 cleavage site may be introduced at the junction of the fusion moiety so that the desired polypeptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith *et al.* (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse
30 glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli*

expression vectors include pTrc (Amann *et al.* (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.* (1990) *Gene Expression Technology: Methods in Enzymology* 185:60-89).

Recombinant protein expression can be maximized in a host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S. (1990) *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California 119-128). Alternatively, the sequence of the polynucleotide of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.* (1992) *Nucleic Acids Res.* 20:2111-2118).

Sulfatase polynucleotides can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari *et al.* (1987) *EMBO J.* 6:229-234), pMFa (Kurjan *et al.* (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.* (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA).

Sulfatase polynucleotides can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow *et al.* (1989) *Virology* 170:31-39).

In certain embodiments of the invention, the polynucleotides described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express sulfatase polynucleotides. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the polynucleotides described herein. These are found for example in Sambrook *et al.*

(1989) *Molecular Cloning: A Laboratory Manual 2nd, ed., Cold Spring Harbor Laboratory*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a

regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the polynucleotide sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to
5 expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such
10 as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection,
15 electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

Host cells can contain more than one vector. Thus, different nucleotide
20 sequences can be introduced on different vectors of the same cell. Similarly, sulfatase polynucleotides can be introduced either alone or with other polynucleotides that are not related to sulfatase polynucleotides such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the sulfatase polynucleotide
25 vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that
30 complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can

be contained in the same vector that contains the polynucleotides described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the polypeptide is desired, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the sulfatase polypeptides or heterologous to these polypeptides.

Where the polypeptide is not secreted into the medium, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The polypeptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the polypeptides described herein, the polypeptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the polypeptides may include an initial modified methionine in some cases as a result of a host-mediated process.

Uses of Vectors and Host Cells

It is understood that "host cells" and "recombinant host cells" refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell.

Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

The host cells expressing the polypeptides described herein, and particularly recombinant host cells, have a variety of uses. First, the cells are useful for producing sulfatase proteins or polypeptides that can be further purified to produce desired amounts of sulfatase protein or fragments. Thus, host cells containing expression vectors are
5 useful for polypeptide production.

Host cells are also useful for conducting cell-based assays involving sulfatase or sulfatase fragments. Thus, a recombinant host cell expressing a native sulfatase is useful to assay for compounds that stimulate or inhibit sulfatase function, gene expression at the level of transcription or translation, and interaction with other cellular components.

10 Host cells are also useful for identifying sulfatase mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant sulfatase (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native sulfatase.

15 Recombinant host cells are also useful for expressing the chimeric polypeptides described herein to assess compounds that activate or suppress activation by means of a heterologous domain, segment, site, and the like, as disclosed herein.

Further, mutant sulfatases can be designed in which one or more of the various functions is engineered to be increased or decreased and used to augment or replace
20 sulfatase proteins in an individual. Thus, host cells can provide a therapeutic benefit by replacing an aberrant sulfatase or providing an aberrant sulfatase that provides a therapeutic result. In one embodiment, the cells provide sulfatases that are abnormally active.

In another embodiment, the cells provide sulfatases that are abnormally inactive.
25 These sulfatases can compete with endogenous sulfatases in the individual.

In another embodiment, cells expressing sulfatases that cannot be activated, are introduced into an individual in order to compete with endogenous sulfatases for substrate. For example, in the case in which excessive substrate or substrate analog is part of a treatment modality, it may be necessary to effectively inactivate the substrate or
30 substrate analog at a specific point in treatment. Providing cells that compete for the molecule, but which cannot be affected by sulfatase activation would be beneficial.

Homologously recombinant host cells can also be produced that allow the *in situ* alteration of endogenous sulfatase polynucleotide sequences in a host cell genome. The host cell includes, but is not limited to, a stable cell line, cell *in vivo*, or cloned microorganism. This technology is more fully described in WO 93/09222, WO 91/12650, WO 91/06667, U.S. 5,272,071, and U.S. 5,641,670. Briefly, specific polynucleotide sequences corresponding to the sulfatase polynucleotides or sequences proximal or distal to a sulfatase gene are allowed to integrate into a host cell genome by homologous recombination where expression of the gene can be affected. In one embodiment, regulatory sequences are introduced that either increase or decrease expression of an endogenous sequence. Accordingly, a sulfatase protein can be produced in a cell not normally producing it. Alternatively, increased expression of sulfatase protein can be effected in a cell normally producing the protein at a specific level. Further, expression can be decreased or eliminated by introducing a specific regulatory sequence. The regulatory sequence can be heterologous to the sulfatase protein sequence or can be a homologous sequence with a desired mutation that affects expression. Alternatively, the entire gene can be deleted. The regulatory sequence can be specific to the host cell or capable of functioning in more than one cell type. Still further, specific mutations can be introduced into any desired region of the gene to produce mutant sulfatase proteins. Such mutations could be introduced, for example, into the specific functional regions such as the peptide substrate-binding site.

In one embodiment, the host cell can be a fertilized oocyte or embryonic stem cell that can be used to produce a transgenic animal containing the altered sulfatase gene. Alternatively, the host cell can be a stem cell or other early tissue precursor that gives rise to a specific subset of cells and can be used to produce transgenic tissues in an animal. See also Thomas *et al.*, *Cell* 51:503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced gene has homologously recombined with the endogenous sulfatase gene is selected (see e.g., Li, E. *et al.* (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras (see e.g., Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant

female foster animal and the embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination
5 vectors and homologous recombinant animals are described further in Bradley, A. (1991) *Current Opinions in Biotechnology* 2:823-829 and in PCT International Publication Nos. WO 90/11354; WO 91/01140; and WO 93/04169.

The genetically engineered host cells can be used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent,
10 such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a sulfatase protein and identifying and
15 evaluating modulators of sulfatase protein activity.

Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

In one embodiment, a host cell is a fertilized oocyte or an embryonic stem cell into which sulfatase polynucleotide sequences have been introduced.

20 A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the sulfatase nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

25 Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the sulfatase protein to particular cells.

30 Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating*

the Mouse Embryo, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A
5 transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

10 In another embodiment, transgenic non-human animals can be produced which contain selected systems, which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al.* (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase
15 system of *S. cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected
20 protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut *et al.* (1997) *Nature* 385:810-813 and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit
25 the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to a pseudopregnant female foster animal. The offspring born of this female animal will be a clone of the animal
30 from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the polypeptides described herein are useful to conduct the assays described herein in an *in vivo* context.

Accordingly, the various physiological factors that are present *in vivo* and that could affect binding or activation, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* sulfatase function, including peptide interaction, the effect of specific mutant sulfatases on sulfatase function and peptide interaction, and the effect of chimeric sulfatases. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more sulfatase functions.

In general, methods for producing transgenic animals include introducing a nucleic acid sequence according to the present invention, the nucleic acid sequence capable of expressing the protein in a transgenic animal, into a cell in culture or *in vivo*. When introduced *in vivo*, the nucleic acid is introduced into an intact organism such that one or more cell types and, accordingly, one or more tissue types, express the nucleic acid encoding the protein. Alternatively, the nucleic acid can be introduced into virtually all cells in an organism by transfecting a cell in culture, such as an embryonic stem cell, as described herein for the production of transgenic animals, and this cell can be used to produce an entire transgenic organism. As described, in a further embodiment, the host cell can be a fertilized oocyte. Such cells are then allowed to develop in a female foster animal to produce the transgenic organism.

Pharmaceutical Compositions

Sulfatase nucleic acid molecules, proteins, modulators of the protein, and antibodies (also referred to herein as "active compounds") can be incorporated into pharmaceutical compositions suitable for administration to a subject, e.g., a human. Such compositions typically comprise the nucleic acid molecule, protein, modulator, or antibody and a pharmaceutically acceptable carrier.

The term "administer" is used in its broadest sense and includes any method of introducing the compositions of the present invention into a subject. This includes producing polypeptides or polynucleotides *in vivo* by *in vivo* transcription or translation of polynucleotides that have been exogenously introduced into a subject. Thus, polypeptides or nucleic acids produced in the subject from the exogenous compositions are encompassed in the term "administer."

As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, such media can be used in the compositions of the invention. Supplementary active compounds can also be incorporated into the compositions. A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can

be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid,

5 thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

10 Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a sulfatase protein or anti-sulfatase antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic
15 dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

20 Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For oral administration, the agent can be contained in enteric forms to survive the stomach or further coated or mixed to be released in a particular region of the GI tract by known methods. For the purpose of oral therapeutic administration, the active compound can be incorporated
25 with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain
30 any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant

such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser, which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound

calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (U.S. 5,328,470) or by stereotactic injection (see e.g., Chen *et al.* (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

As defined herein, a therapeutically effective amount of protein or polypeptide (i.e., an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight.

The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody, protein, or polypeptide in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody, protein, or polypeptide used for treatment may increase or decrease over the course of a particular treatment. Changes

in dosage may result and become apparent from the results of diagnostic assays as described herein.

The present invention encompasses agents which modulate expression or activity. An agent may, for example, be a small molecule. For example, such small
5 molecules include, but are not limited to, peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about
10 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

It is understood that appropriate doses of small molecule agents depends upon
15 a number of factors within the ken of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of the small molecule will vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the small molecule to have
20 upon the nucleic acid or polypeptide of the invention. Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is
25 furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. Such appropriate doses may be determined using the assays described herein. When one or more of these small molecules is to be administered to an animal (e.g., a human) in order to modulate expression or activity of a polypeptide or nucleic
30 acid of the invention, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose

level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

This invention may be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will fully convey the invention to those skilled in the art. Many modifications and other embodiments of the invention will come to mind in one skilled in the art to which this invention pertains having the benefit of the teachings presented in the foregoing description. Although specific terms are employed, they are used as in the art unless otherwise indicated.

THAT WHICH IS CLAIMED:

1 An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5
- (a) the amino acid sequence shown in SEQ ID NO:1;
 - (b) the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
 - (c) the amino acid sequence of an allelic variant of the amino acid sequence shown in SEQ ID NO:1;
 - 10 (d) the amino acid sequence of an allelic variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
 - (e) the amino acid sequence of a sequence variant of the amino acid sequence shown in SEQ ID NO:1, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing to the nucleic acid molecule shown in SEQ ID NO:2 under
 - 15 stringent conditions;
 - (f) the amino acid sequence of a sequence variant of the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. _____, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing under stringent conditions to the cDNA contained in ATCC Deposit No. _____;
 - 20 (g) a fragment of the amino acid sequence shown in SEQ ID NO:1, said fragment having sulfatase activity;
 - (h) a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment has sulfatase activity;
 - (i) the amino acid sequence of the mature sulfatase polypeptide,
 - 25 comprising about amino acid residues 17 to the last amino acid of SEQ ID NO:1;
 - (j) the amino acid sequence of the mature polypeptide, comprising from about amino acid 17 to the last amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;
 - (k) the amino acid sequence of an epitope bearing region of a
 - 30 polypeptide of (a)-(j).

2. An isolated antibody that selectively binds to a polypeptide of claim 1.

3. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 5
- (a) the nucleotide sequence shown in SEQ ID NO:2;
 - (b) the nucleotide sequence in the cDNA contained in ATCC Deposit No. ____;
 - (c) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:1;
 - 10 (d) a nucleotide sequence encoding the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____;
 - (e) a nucleotide sequence encoding a polypeptide of claim 1; and
 - (f) a nucleotide sequence complementary to a nucleotide sequence of (a)-(e).
- 15

4. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 20
- (a) a nucleotide sequence encoding a variant of the amino acid sequence shown in SEQ ID NO:1, wherein said nucleotide sequence hybridizes to the nucleotide sequence shown in SEQ ID NO:2 under stringent conditions;
 - (b) a nucleotide sequence encoding a variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____, wherein the nucleic acid sequence of the sequence variant hybridizes to the cDNA contained in ATCC Deposit No. ____ under stringent conditions;
 - 25 (c) a nucleotide sequence encoding the polypeptide of claim 2; and
 - (d) a nucleotide sequence complementary to either of the nucleotide sequences in (a)-(c).

5. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

- 30
- (a) a nucleotide sequence encoding a fragment of the amino acid sequence shown in SEQ ID NO:1, wherein the fragment encodes a sulfatase activity;

A

(b) a nucleotide sequence encoding a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment encodes a sulfatase activity;

(c) a nucleotide sequence encoding residues 17 to the last amino acid shown in SEQ ID NO:1;

(d) a nucleotide sequence encoding residues 17 to the last amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(e) a nucleotide sequence complementary to a nucleotide sequences of (a)-(d).

6. A nucleic acid vector comprising the nucleic acid sequences in any of claims 3-5.

7. A host cell containing the vector of claim 6.

8. A method for producing a sulfatase polypeptide, comprising: introducing a nucleotide sequence encoding a polypeptide sequence of claim 1 into a host cell, and culturing the host cell under conditions in which said polypeptide is expressed.

9. A method for detecting the presence of any of the polypeptides in claim 1 in a sample, said method comprising contacting said sample with an agent that specifically allows detection of the presence of the polypeptide in the sample and then detecting the presence of the polypeptide.

10. The method of claim 9, wherein said agent is capable of selective physical association with said polypeptide.

11. The method of claim 10, wherein said agent binds to said polypeptide.

12. The method of claim 11, wherein said agent is an antibody.

13. The method of claim 11, wherein said agent is a sulfated conjugate.

14. A kit comprising reagents used for the method of claim 9, wherein the reagents comprise an agent that specifically binds to said polypeptide.

15. A method for detecting the presence of any of the nucleic acid molecules in any of claims 3-5 in a sample, the method comprising contacting said sample with an agent that specifically allows detection of the presence of the nucleic acid molecule in the sample and then detecting the presence of the nucleic acid molecule.

16. The method of claim 15, wherein said method comprises contacting the sample with an oligonucleotide that hybridizes to the nucleic acid sequences under stringent conditions and determining whether the oligonucleotide binds to the nucleic acid sequence in the sample.

17. The method of claim 15, wherein the nucleic acid, whose presence is detected, is mRNA.

18. A kit comprising reagents used for the method of claim 15, wherein the reagents comprise a compound that hybridizes under stringent conditions to any of the nucleic acid molecules.

19. A method for identifying an agent that interacts with any of the polypeptides of claim 1 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent and measuring the interaction.

20. A method of screening a cell to identify an agent that interacts with any of the polypeptides of claim 1 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent, and measuring the interaction.

21. A method for identifying an agent that binds to any of the polypeptides in claim 1, said method comprising contacting the polypeptide with an agent that binds to the polypeptide and assaying the complex formed with the agent bound to the polypeptide.

22. The method of claim 21, wherein a fragment of the polypeptide is contacted.

23. A method of screening a cell to identify an agent that modulates the level or activity of any of the polypeptides of claim 1 in a cell, said method comprising: contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

24. The method of claim 19 wherein said cell is a cell from a DRG neuron, spinal cord, or brain.

25. The method of claim 23 wherein said agent increases the level or activity of said polypeptide.

26. The method of claim 23 wherein said agent decreases the level or activity of said polypeptide.

27. The method of claim 19, said method comprising: (1) exposing said agent to said polypeptide under conditions that allow said agent to interact with said polypeptide; (2) adding competing polypeptide that can interact with said agent; and (3) comparing the amount of interaction between said agent and said polypeptide to the amount of interaction in the absence of said competing polypeptide.

28. The method of claim 19 wherein said interaction is binding

29. The method of claim 23 wherein said agent increases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

30. The method of claim 23 wherein said agent decreases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

31. The method of claim 23 wherein said cell is *in vivo*.

32. The method of claim 31 wherein said cell is in a transgenic animal.

33. The method of claim 31 wherein said cell is in a non-transgenic subject.

34. The method of claim 23 wherein said cell is *in vitro*.

35. The method of claim 34 wherein said cell has been disrupted.

36. The method of claim 34 wherein said cell is in a biopsy.

37. The method of claim 35 wherein said cell is in cell culture.

38. The method of claim 37 wherein said cell is naturally-occurring or recombinant.

39. The method of claim 23 wherein said agent is selected from the group consisting of a sulfated conjugate; peptide; phosphopeptide; antibody; organic molecule; and inorganic molecule.

40. A method for modulating the level or activity of any of the polypeptides of claim 1, said method comprising contacting said polypeptide with an agent under conditions that allow the agent to modulate the level or activity of the polypeptide.

41. A method for identifying an agent that modulates the level or activity of any of the polypeptides of claim 1 in a cell, said method comprising contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

42. A method for identifying an agent that modulates the level or activity of any of the nucleic acid molecules of claims 3-5 in a cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring said nucleic acid molecule level or activity.

43. A method of screening a cell to identify an agent that modulates the level or activity of any of the nucleic acid molecules in claims 3-5 in said cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring nucleic acid molecule level or activity.

44. A method for identifying an agent that interacts with any of the nucleic acid molecules of claims 3-5 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and

said agent such that said nucleic acid molecule can interact with said agent in measuring the interaction.

45. A method of screening a cell to identify an agent that interacts with any of the nucleic acid molecules of claims 3-5 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent and measuring the interaction.

46. A method for modulating the level or activity of any of the nucleic acid molecules of claims 3-5, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level or activity of the nucleic acid molecule.

47. The method of claim 46 wherein said modulation is in cells derived from tissue selected from the group consisting of DRG neurons, spinal cord, and brain.

48. The method of claim 46 wherein said modulation is *in vivo*.

49. The method of claim 48 wherein said modulation is in a patient having or predisposed to having a disorder involving DRG neurons, spinal cord, or brain.

50. The method of claim 49 wherein said modulation is in a patient having a disorder involving pain.

51. A method of treating a disorder involving DRG neurons, spinal cord, or brain in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 1 to said subject in a therapeutically effective amount.

A

52. A method of treating a disorder involving DRG neurons, spinal cord, or brain in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 1 to said subject in therapeutically effective amounts.

5 53. A pharmaceutical composition containing any of the polypeptides in claim 1 in a pharmaceutically acceptable carrier.

54. A pharmaceutically acceptable composition containing any of the nucleic acid molecules of claims 3-5 in a pharmaceutically acceptable carrier.

10 55. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 3-5.

15 56. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 3-5, wherein said cell expresses any of the polypeptides of claim 1.

20 57. A method for producing a transgenic animal according to claim 46, said method comprising introducing any of the nucleic acid sequences of claims 3-5 into a cell, wherein said cell is present in said animal or gives rise to said animal.

58. An agent identified by any of the methods of claims 19-39

25 59. An agent identified by any of the methods of claims 41-45.

60. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence shown in SEQ ID NO:3;

(b) the amino acid sequence encoded by the cDNA contained in
30 ATCC Deposit No. _____;

(c) the amino acid sequence of an allelic variant of the amino acid sequence shown in SEQ ID NO:3;

(d) the amino acid sequence of an allelic variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;

(e) the amino acid sequence of a sequence variant of the amino acid sequence shown in SEQ ID NO:3, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing to the nucleic acid molecule shown in SEQ ID NO:4 under stringent conditions;

(f) the amino acid sequence of a sequence variant of the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. _____, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing under stringent conditions to the cDNA contained in ATCC Deposit No. _____;

(g) a fragment of the amino acid sequence shown in SEQ ID NO:3, wherein the fragment comprises at least 6 contiguous amino acids from nucleotides 510-670;

(h) a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 6 contiguous amino acids from nucleotides 510-670;

(i) the amino acid sequence of the mature sulfatase polypeptide, comprising about amino acid residues 23-871 of SEQ ID NO:3;

(j) the amino acid sequence of the mature polypeptide, comprising from about amino acid 23 to the last amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(k) the amino acid sequence of an epitope bearing region of a polypeptide of (a)-(j).

61. An isolated polypeptide having at least 94% or more sequence identity with the amino acid sequence of SEQ ID NO:3.

62. The polypeptide a claim 61, wherein said polypeptide has at least 97% or more sequence identity with the amino acid sequence of SEQ ID NO:3.

63. The polypeptide a claim 61, wherein said polypeptide has at least 98% or more sequence identity with the amino acid sequence of SEQ ID NO:3.

64. The polypeptide a claim 61, wherein said polypeptide has at least 99% or more sequence identity with the amino acid sequence of SEQ ID NO:3.

5 65. An isolated antibody that selectively binds to a polypeptide of claim 60.

66. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 10 (a) the nucleotide sequence shown in SEQ ID NO:4;
(b) the nucleotide sequence in the cDNA contained in ATCC Deposit No. ____;
(c) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:3;
(d) a nucleotide sequence encoding the amino acid sequence encoded
15 by the cDNA contained in ATCC Deposit No. ____;
(e) a nucleotide sequence encoding a polypeptide of claim 60; and
(f) a nucleotide sequence complementary to a nucleotide sequence of (a)-(e).

20 67. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a variant of the amino acid sequence shown in SEQ ID NO:3, wherein said nucleotide sequence hybridizes to the nucleotide sequence shown in SEQ ID NO:4 under stringent conditions;
25 (b) a nucleotide sequence encoding a variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____, wherein the nucleic acid sequence of the sequence variant hybridizes to the cDNA contained in ATCC Deposit No. ____ under stringent conditions;
(c) a nucleotide sequence encoding the polypeptide of claim 61; and
30 (d) a nucleotide sequence complementary to either of the nucleotide sequences in (a)-(c).

68. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a fragment of the amino acid sequence shown in SEQ ID NO:3, wherein the fragment comprises at least 6 contiguous amino acids from nucleotides 510-670;

(b) a nucleotide sequence encoding a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 6 contiguous amino acids from nucleotides 510-670;

(c) a nucleotide sequence comprising at least 18 consecutive nucleotides of the sequence shown in SEQ ID NO:4 from 1-670, 3008-3514, and 3998-4321;

(d) a nucleotide sequence comprising at least 18 consecutive nucleotides from 1-670, 3008-3514, and 3998-4321 of SEQ ID NO:4;

(e) a nucleotide sequence encoding residues 23-871 of the amino acid shown in SEQ ID NO:3;

(f) a nucleotide sequence encoding residues 23-871 of the amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(g) a nucleotide sequence complementary to a nucleotide sequence of (a)-(f).

69. A nucleic acid vector comprising a nucleic acid sequence in any of claims 66-68.

70. A host cell containing the vector of claim 69.

71. A method for producing a sulfatase polypeptide, comprising: introducing a nucleotide sequence encoding a polypeptide sequence of claim 60 into a host cell, and culturing the host cell under conditions in which said polypeptide is expressed.

72. A method for detecting the presence of any of the polypeptides in claim 60 in a sample, said method comprising contacting said sample with an agent

that specifically allows detection of the presence of the polypeptide in the sample and then detecting the presence of the polypeptide.

5 73. The method of claim 72, wherein said agent is capable of selective physical association with said polypeptide.

74. The method of claim 73, wherein said agent binds to said polypeptide.

10 75. The method of claim 74, wherein said agent is an antibody.

76. The method of claim 74, wherein said agent is a peptide substrate.

15 77. A kit comprising reagents used for the method of claim 72, wherein the reagents comprise an agent that specifically binds to said polypeptide.

20 78. A method for detecting the presence of any of the nucleic acid molecules in any of claims 66-68 in a sample, the method comprising contacting said sample with an agent that specifically allows detection of the presence of the nucleic acid molecule in the sample and then detecting the presence of the nucleic acid molecule.

25 79. The method of claim 78, wherein said method comprises contacting the sample with an oligonucleotide that hybridizes to the nucleic acid sequences under stringent conditions and determining whether the oligonucleotide binds to the nucleic acid sequence in the sample.

80. The method of claim 78, wherein the nucleic acid, whose presence is detected, is mRNA.

30 81. A kit comprising reagents used for the method of claim 78, wherein the reagents comprise a compound that hybridizes under stringent conditions to any of the nucleic acid molecules.

A

82. A method for identifying an agent that interacts with any of the polypeptides of claim 60 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent and measuring the interaction.

83. A method of screening a cell to identify an agent that interacts with any of the polypeptides of claim 60 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent, and measuring the interaction.

84. A method for identifying an agent that binds to any of the polypeptides in claim 60, said method comprising contacting the polypeptide with an agent that binds to the polypeptide and assaying the complex formed with the agent bound to the polypeptide.

85. The method of claim 84, wherein a fragment of the polypeptide is contacted.

86. A method of screening a cell to identify an agent that modulates the level or activity of any of the polypeptides of claim 60 in a cell, said method comprising: contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

87. The method of claim 86 wherein said cell is a cell from a breast or colon carcinoma.

88. The method of claim 86 wherein said agent increases the level or activity of said polypeptide.

89. The method of claim 86 wherein said agent decreases the level or activity of said polypeptide.

90. The method of claim 86, said method comprising: (1) exposing said agent to said polypeptide under conditions that allow said agent to interact with said polypeptide; (2) adding competing polypeptide that can interact with said agent; and (3) comparing the amount of interaction between said agent and said polypeptide to the amount of interaction in the absence of said competing polypeptide.

91. The method of claim 82 wherein said interaction is binding.

92. The method of claim 86 wherein said agent increases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

93. The method of claim 86 wherein said agent decreases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

94. The method of claim 86 wherein said cell is *in vivo*.

95. The method of claim 94 wherein said cell is in a transgenic animal.

96. The method of claim 94 wherein said cell is in a non-transgenic subject.

- 5
97. The method of claim 86 wherein said cell is *in vitro*.
98. The method of claim 97 wherein said cell has been disrupted.
99. The method of claim 97 wherein said cell is in a biopsy.
100. The method of claim 98 wherein said cell is in cell culture.
101. The method of claim 100 wherein said cell is naturally-occurring or
10 recombinant.
102. The method of claim 86 wherein said agent is selected from the group
consisting of a peptide; phosphopeptide; antibody; organic molecule; and inorganic
molecule.
- 15
103. A method for modulating the level or activity of any of the
polypeptides of claim 60, said method comprising contacting said polypeptide with an
agent under conditions that allow the agent to modulate the level or activity of the
polypeptide.
- 20
104. A method for identifying an agent that modulates the level or activity
of any of the polypeptides of claim 60 in a cell, said method comprising contacting
said agent with a cell capable of expressing said polypeptide such that said
polypeptide level or activity can be modulated in said cell by said agent and
25 measuring said polypeptide level or activity.
105. A method for identifying an agent that modulates the level or activity
of any of the nucleic acid molecules of claims 66-68 in a cell, said method comprising
contacting said agent with the cell capable of expressing said nucleic acid molecule
30 such that said nucleic acid molecule level or activity can be modulated in said cell by
said agent and measuring said nucleic acid molecule level or activity.

106. A method of screening a cell to identify an agent that modulates the level or activity of any of the nucleic acid molecules in claims 66-68 in said cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring nucleic acid molecule level or activity.

107. A method for identifying an agent that interacts with any of the nucleic acid molecules of claims 66-68 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent in measuring the interaction.

108. A method of screening a cell to identify an agent that interacts with any of the nucleic acid molecules of claims 66-68 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent and measuring the interaction.

109. A method for modulating the level or activity of any of the nucleic acid molecules of claims 66-68, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level or activity of the nucleic acid molecule.

110. The method of claim 109 wherein said modulation is in cells derived from tissue selected from the group consisting of breast and colon carcinoma.

111. The method of claim 109 wherein said modulation is *in vivo*.

112. The method of claim 111 wherein said modulation is in a patient having a disorder involving the breast or colon, or predisposed to having breast or colon carcinoma.

113. The method of claim 112 wherein said modulation is in a patient having or predisposed to having breast or colon carcinoma.

5 114. A method of treating a disorder involving breast or colon in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 60 to said subject in a therapeutically effective amount.

10 115. A method of treating breast or colon carcinoma in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 60 to said subject in therapeutically effective amounts.

15 116. A pharmaceutical composition containing any of the polypeptides in claim 60 in a pharmaceutically acceptable carrier.

117. A pharmaceutically acceptable composition containing any of the nucleic acid molecules of claims 66-68 in a pharmaceutically acceptable carrier.

20 118. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 66-68.

25 119. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 66-68, wherein said cell expresses any of the polypeptides of claim 60.

120. A method for producing a transgenic animal according to claim 119, said method comprising introducing any of the nucleic acid sequences of claims 66-68 into a cell, wherein said cell is present in said animal or gives rise to said animal.

30 121. An agent identified by any of the methods of claims 82-102.

122. An agent identified by any of the methods of claims 104-108.

123. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5
- (a) the amino acid sequence shown in SEQ ID NO:5;
 - (b) the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
 - (c) the amino acid sequence of an allelic variant of the amino acid sequence shown in SEQ ID NO:5;
 - 10 (d) the amino acid sequence of an allelic variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
 - (e) the amino acid sequence of a sequence variant of the amino acid sequence shown in SEQ ID NO:5, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing to the nucleic acid molecule shown in SEQ ID NO:6 under stringent conditions;
 - 15 (f) the amino acid sequence of a sequence variant of the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. _____, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing under stringent conditions to the cDNA contained in ATCC Deposit No. _____;
 - (g) a fragment of the amino acid sequence shown in SEQ ID NO:5,
 - 20 wherein the fragment comprises at least 22 contiguous amino acids;
 - (h) a fragment of the amino acid sequence shown in SEQ ID NO:5, wherein said fragment comprises at least 22 contiguous amino acids.
 - (i) a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 22
 - 25 contiguous amino acids;
 - (j) the amino acid sequence of the mature sulfatase polypeptide, comprising about amino acid residues 24-569 of SEQ ID NO:5;
 - (k) the amino acid sequence of the mature polypeptide, comprising from about amino acid 24 to the last amino acid encoded by the cDNA clone contained
 - 30 in ATCC Deposit No. _____;
 - (l) the amino acid sequence of an epitope bearing region of a polypeptide of (a)-(k).

124. An isolated polypeptide having at least 50% sequence identity with the amino acid sequence of SEQ ID NO:5.

125. The polypeptide a claim 124, wherein said polypeptide has at least 75% sequence identity with the amino acid sequence of SEQ ID NO:5.

126. The polypeptide a claim 124, wherein said polypeptide has at least 85% sequence identity with the amino acid sequence of SEQ ID NO:5.

127. The polypeptide a claim 124, wherein said polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO:5.

128. An isolated antibody that selectively binds to a polypeptide of claim 123.

129. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence shown in SEQ ID NO:6;
- (b) the nucleotide sequence in the cDNA contained in ATCC Deposit No. ____;
- (c) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:5;
- (d) a nucleotide sequence encoding the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____;
- (e) a nucleotide sequence encoding a polypeptide of claim 123; and
- (f) a nucleotide sequence complementary to a nucleotide sequence of (a)-(e).

130. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a variant of the amino acid sequence shown in SEQ ID NO:5, wherein said nucleotide sequence hybridizes to the nucleotide sequence shown in SEQ ID NO:6 under stringent conditions;

(b) a nucleotide sequence encoding a variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the nucleic acid sequence of the sequence variant hybridizes to the cDNA contained in ATCC Deposit No. _____ under stringent conditions;

(c) a nucleotide sequence encoding the polypeptide of claim 124; and

(d) a nucleotide sequence complementary to either of the nucleotide sequences in (a)-(c).

131. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a fragment of the amino acid sequence shown in SEQ ID NO:5, wherein the fragment comprises at least 22 contiguous amino acids;

(b) a nucleotide sequence encoding a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 22 contiguous amino acids;

(c) a nucleotide sequence comprising at least 24 consecutive nucleotides of the sequence shown in SEQ ID NO:6 from 30-1900 and 1930-2420;

(d) a nucleotide sequence comprising at least 24 consecutive nucleotides from 30-1900 and 1930-2420 of SEQ ID NO:6;

(e) a nucleotide sequence encoding residues 24-569 of the amino acid shown in SEQ ID NO:5;

(f) a nucleotide sequence encoding residues 24-569 of the amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(g) a nucleotide sequence complementary to a nucleotide sequences of (a)-(f).

132. A nucleic acid vector comprising the nucleic acid sequences in any of claims 129-131.

133. A host cell containing the vector of claim 132.

134. A method for producing a sulfatase polypeptide, comprising: introducing a nucleotide sequence encoding a polypeptide sequence of claim 123 into a host cell, and culturing the host cell under conditions in which said polypeptide is expressed.

135. A method for detecting the presence of any of the polypeptides in claim 123 in a sample, said method comprising contacting said sample with an agent that specifically allows detection of the presence of the polypeptide in the sample and then detecting the presence of the polypeptide.

136. The method of claim 135, wherein said agent is capable of selective physical association with said polypeptide.

137. The method of claim 136, wherein said agent binds to said polypeptide.

138. The method of claim 137, wherein said agent is an antibody.

139. The method of claim 137, wherein said agent is a sulfated conjugate substrate.

140. A kit comprising reagents used for the method of claim 135, wherein the reagents comprise an agent that specifically binds to said polypeptide.

141. A method for detecting the presence of any of the nucleic acid molecules in any of claims 129-131 in a sample, the method comprising contacting said sample with an agent that specifically allows detection of the presence of the

X
nucleic acid molecule in the sample and then detecting the presence of the nucleic acid molecule.

5 142. The method of claim 141, wherein said method comprises contacting the sample with an oligonucleotide that hybridizes to the nucleic acid sequences under stringent conditions and determining whether the oligonucleotide binds to the nucleic acid sequence in the sample.

10 143. The method of claim 141, wherein the nucleic acid, whose presence is detected, is mRNA.

15 144. A kit comprising reagents used for the method of claim 141, wherein the reagents comprise a compound that hybridizes under stringent conditions to any of the nucleic acid molecules.

20 145. A method for identifying an agent that interacts with any of the polypeptides of claim 123 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent and measuring the interaction.

25 146. A method of screening a cell to identify an agent that interacts with any of the polypeptides of claim 123 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent, and measuring the interaction.

30 147. A method for identifying an agent that binds to any of the polypeptides in claim 123, said method comprising contacting the polypeptide with an agent that binds to the polypeptide and assaying the complex formed with the agent bound to the polypeptide.

A
148. The method of claim 147, wherein a fragment of the polypeptide is contacted.

149. A method of screening a cell to identify an agent that modulates the level or activity of any of the polypeptides of claim 123 in a cell, said method comprising: contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

150. The method of claim 149 wherein said cell is a cell from a colon carcinoma.

151. The method of claim 149 wherein said agent increases the level or activity of said polypeptide.

152. The method of claim 149 wherein said agent decreases the level or activity of said polypeptide.

153. The method of claim 145, said method comprising: (1) exposing said agent to said polypeptide under conditions that allow said agent to interact with said polypeptide; (2) adding competing polypeptide that can interact with said agent; and (3) comparing the amount of interaction between said agent and said polypeptide to the amount of interaction in the absence of said competing polypeptide.

154. The method of claim 153 wherein said interaction is binding.

155. The method of claim 149 wherein said agent increases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

156. The method of claim 149 wherein said agent decreases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

157. The method of claim 149 wherein said cell is *in vivo*.

158. The method of claim 157 wherein said cell is in a transgenic animal.

159. The method of claim 157 wherein said cell is in a non-transgenic subject.

160. The method of claim 149 wherein said cell is *in vitro*.

161. The method of claim 160 wherein said cell has been disrupted.

162. The method of claim 160 wherein said cell is in a biopsy.

163. The method of claim 161 wherein said cell is in cell culture.

164. The method of claim 163 wherein said cell is naturally-occurring or recombinant.

165. The method of claim 149 wherein said agent is selected from the group consisting of a sulfated conjugate; peptide; phosphopeptide; antibody; organic molecule; and inorganic molecule.

166. A method for modulating the level or activity of any of the polypeptides of claim 123, said method comprising contacting said polypeptide with

A
an agent under conditions that allow the agent to modulate the level or activity of the polypeptide.

5 167. A method for identifying an agent that modulates the level or activity of any of the polypeptides of claim 123 in a cell, said method comprising contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

10 168. A method for identifying an agent that modulates the level or activity of any of the nucleic acid molecules of claims 129-131 in a cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring said nucleic acid molecule level or activity.

15 169. A method of screening a cell to identify an agent that modulates the level or activity of any of the nucleic acid molecules in claims 129-131 in said cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be
20 modulated in said cell by said agent and measuring nucleic acid molecule level or activity.

25 170. A method for identifying an agent that interacts with any of the nucleic acid molecules of claims 129-131 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent in measuring the interaction.

30 171. A method of screening a cell to identify an agent that interacts with any of the nucleic acid molecules of claims 129-131 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said

nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent and measuring the interaction.

5 172. A method for modulating the level or activity of any of the nucleic acid molecules of claims 129-131, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level or activity of the nucleic acid molecule.

10 173. The method of claim 172 wherein said modulation is in cells derived from colon tissue.

174. The method of claim 172 wherein said modulation is *in vivo*.

15 175. The method of claim 174 wherein said modulation is in a patient having a disorder involving the colon.

176. The method of claim 175 wherein said modulation is in a patient having colon carcinoma.

20 177. A method of treating a disorder involving the colon in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 123 to said subject in a therapeutically effective amount.

25 178. A method of treating colon carcinoma in a subject in need of such treatment, said method comprising administering any of the polypeptides of claim 123 to said subject in therapeutically effective amounts.

30 179. A pharmaceutical composition containing any of the polypeptides in claim 123 in a pharmaceutically acceptable carrier.

180. A pharmaceutically acceptable composition containing any of the nucleic acid molecules of claims 129-131 in a pharmaceutically acceptable carrier.

181. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 129-131.

5 182. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 129-131, wherein said cell expresses any of the polypeptides of claim 123.

10 183. A method for producing a transgenic animal according to claim 182, said method comprising introducing any of the nucleic acid sequences of claims 129-131 into a cell, wherein said cell is present in said animal or gives rise to said animal.

184. An agent identified by any of the methods of claims 145-165.

15 185. An agent identified by any of the methods of claims 167-171.

186. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence shown in SEQ ID NO:7;
- (b) the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
- (c) the amino acid sequence of an allelic variant of the amino acid sequence shown in SEQ ID NO:7;
- 25 (d) the amino acid sequence of an allelic variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____;
- (e) the amino acid sequence of a sequence variant of the amino acid sequence shown in SEQ ID NO:7, wherein the sequence variant is encoded by a nucleic acid molecule hybridizing to the nucleic acid molecule shown in SEQ ID NO:8 under stringent conditions;
- 30 (f) the amino acid sequence of a sequence variant of the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. _____, wherein the

sequence variant is encoded by a nucleic acid molecule hybridizing under stringent conditions to the cDNA contained in ATCC Deposit No. _____;

(g) a fragment of the amino acid sequence shown in SEQ ID NO:7, wherein the fragment comprises at least 11 contiguous amino acids;

(h) a fragment of the amino acid sequence shown in SEQ ID NO:7, wherein said fragment comprises at least 11 contiguous amino acids.

(i) a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 11 contiguous amino acids;

(j) the amino acid sequence of the mature sulfatase polypeptide, comprising about amino acid residues 50-550 of SEQ ID NO:7;

(k) the amino acid sequence of the mature polypeptide, comprising from about amino acid 50 to the last amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(l) the amino acid sequence of an epitope bearing region of a polypeptide of (a)-(k).

187. An isolated polypeptide having at least 50% sequence identity with the amino acid sequence of SEQ ID NO:7.

188. The polypeptide a claim 187, wherein said polypeptide has at least 75% sequence identity with the amino acid sequence of SEQ ID NO:7.

189. The polypeptide a claim 187, wherein said polypeptide has at least 85% sequence identity with the amino acid sequence of SEQ ID NO:7.

190. The polypeptide a claim 187, wherein said polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO:7.

191. An isolated antibody that selectively binds to a polypeptide of claim 186.

192. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 5
- (a) the nucleotide sequence shown in SEQ ID NO:8;
- (b) the nucleotide sequence in the cDNA contained in ATCC Deposit No. ____;
- (c) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:7;
- (d) a nucleotide sequence encoding the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____;
- 10 (e) a nucleotide sequence encoding a polypeptide of claim 186; and
- (f) a nucleotide sequence complementary to a nucleotide sequence of (a)-(e).

193. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) a nucleotide sequence encoding a variant of the amino acid sequence shown in SEQ ID NO:7, wherein said nucleotide sequence hybridizes to the nucleotide sequence shown in SEQ ID NO:8 under stringent conditions;
- (b) a nucleotide sequence encoding a variant of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. ____, wherein the nucleic acid sequence of the sequence variant hybridizes to the cDNA contained in ATCC Deposit No. ____ under stringent conditions;
- 20 (c) a nucleotide sequence encoding the polypeptide of claim 187; and
- 25 (d) a nucleotide sequence complementary to either of the nucleotide sequences in (a)-(c).

194. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

- 30 (a) a nucleotide sequence encoding a fragment of the amino acid sequence shown in SEQ ID NO:7, wherein the fragment comprises at least 11 contiguous amino acids;

(b) a nucleotide sequence encoding a fragment of the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. _____, wherein the fragment comprises at least 11 contiguous amino acids;

5 (c) a nucleotide sequence comprising at least 17 consecutive nucleotides of the sequence shown in SEQ ID NO:8 from 272-751, 1074-1264, and 20 from 1264-1551;

(d) a nucleotide sequence comprising at least 17 consecutive nucleotides from 272-751, 1074-1264, and 20 from 1264-1551 of SEQ ID NO:8;

10 (e) a nucleotide sequence encoding residues 50-550 of the amino acid shown in SEQ ID NO:7;

(f) a nucleotide sequence encoding residues 50-550 of the amino acid encoded by the cDNA clone contained in ATCC Deposit No. _____;

(g) a nucleotide sequence complementary to a nucleotide sequences of (a)-(f).

15 195. A nucleic acid vector comprising the nucleic acid sequences in any of claims 193-194.

196. A host cell containing the vector of claim 195.

20 197. A method for producing a sulfatase polypeptide, comprising: introducing a nucleotide sequence encoding a polypeptide sequence of claim 186 into a host cell, and culturing the host cell under conditions in which said polypeptide is expressed.

25 198. A method for detecting the presence of any of the polypeptides in claim 186 in a sample, said method comprising contacting said sample with an agent that specifically allows detection of the presence of the polypeptide in the sample and then detecting the presence of the polypeptide.

30 199. The method of claim 198, wherein said agent is capable of selective physical association with said polypeptide.

200. The method of claim 199, wherein said agent binds to said polypeptide.

201. The method of claim 200, wherein said agent is an antibody.

202. The method of claim 200, wherein said agent is a peptide substrate.

203. A kit comprising reagents used for the method of claim 199, wherein the reagents comprise an agent that specifically binds to said polypeptide.

204. A method for detecting the presence of any of the nucleic acid molecules in any of claims 193-195 in a sample, the method comprising contacting said sample with an agent that specifically allows detection of the presence of the nucleic acid molecule in the sample and then detecting the presence of the nucleic acid molecule.

205. The method of claim 204, wherein said method comprises contacting the sample with an oligonucleotide that hybridizes to the nucleic acid sequences under stringent conditions and determining whether the oligonucleotide binds to the nucleic acid sequence in the sample.

206. The method of claim 204, wherein the nucleic acid, whose presence is detected, is mRNA.

207. A kit comprising reagents used for the method of claim 204, wherein the reagents comprise a compound that hybridizes under stringent conditions to any of the nucleic acid molecules.

208. A method for identifying an agent that interacts with any of the polypeptides of claim 186 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent and measuring the interaction.

209. A method of screening a cell to identify an agent that interacts with any of the polypeptides of claim 186 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said polypeptide and said agent such that said polypeptide can interact with said agent, and measuring the interaction.

210. A method for identifying an agent that binds to any of the polypeptides in claim 186, said method comprising contacting the polypeptide with an agent that binds to the polypeptide and assaying the complex formed with the agent bound to the polypeptide.

211. The method of claim 210, wherein a fragment of the polypeptide is contacted.

212. A method of screening a cell to identify an agent that modulates the level or activity of any of the polypeptides of claim 186 in a cell, said method comprising: contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

213. The method of claim 212 wherein said agent increases the level or activity of said polypeptide.

214. The method of claim 212 wherein said agent decreases the level or activity of said polypeptide.

215. The method of claim 208, said method comprising: (1) exposing said agent to said polypeptide under conditions that allow said agent to interact with said polypeptide; (2) adding competing polypeptide that can interact with said agent; and (3) comparing the amount of interaction between said agent and said polypeptide to the amount of interaction in the absence of said competing polypeptide.

216. The method of claim 208 wherein said interaction is binding.

217. The method of claim 212 wherein said agent increases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

218. The method of claim 212 wherein said agent decreases interaction between said polypeptide and a target molecule for said polypeptide, said method comprising: combining said polypeptide with said agent under conditions that allow said polypeptide to interact with said target molecule; and detecting the formation of a complex between said polypeptide and said target molecule or activity of said polypeptide as a result of interaction of said polypeptide with said target molecule.

219. The method of claim 212 wherein said cell is *in vivo*.

220. The method of claim 219 wherein said cell is in a transgenic animal.

221. The method of claim 219 wherein said cell is in a non-transgenic subject.

222. The method of claim 212 wherein said cell is *in vitro*.

223. The method of claim 222 wherein said cell has been disrupted.

224. The method of claim 222 wherein said cell is in a biopsy.

225. The method of claim 223 wherein said cell is in cell culture.

226. The method of claim 225 wherein said cell is naturally-occurring or recombinant.

227. The method of claim 212 wherein said agent is selected from the group consisting of a sulfated conjugate; peptide; phosphopeptide; antibody; organic molecule; and inorganic molecule.

228. A method for modulating the level or activity of any of the polypeptides of claim 186, said method comprising contacting said polypeptide with an agent under conditions that allow the agent to modulate the level or activity of the polypeptide.

229. A method for identifying an agent that modulates the level or activity of any of the polypeptides of claim 186 in a cell, said method comprising contacting said agent with a cell capable of expressing said polypeptide such that said polypeptide level or activity can be modulated in said cell by said agent and measuring said polypeptide level or activity.

230. A method for identifying an agent that modulates the level or activity of any of the nucleic acid molecules of claims 193-195 in a cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring said nucleic acid molecule level or activity.

231. A method of screening a cell to identify an agent that modulates the level or activity of any of the nucleic acid molecules in claims 193-195 in said cell, said method comprising contacting said agent with the cell capable of expressing said nucleic acid molecule such that said nucleic acid molecule level or activity can be modulated in said cell by said agent and measuring nucleic acid molecule level or activity.

232. A method for identifying an agent that interacts with any of the nucleic acid molecules of claims 193-195 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent in measuring the interaction.

233. A method of screening a cell to identify an agent that interacts with any of the nucleic acid molecules of claims 193-195 in a cell, said method comprising contacting said agent with a cell capable of allowing an interaction between said nucleic acid molecule and said agent such that said nucleic acid molecule can interact with said agent and measuring the interaction.

234. A method for modulating the level or activity of any of the nucleic acid molecules of claims 193-195, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level or activity of the nucleic acid molecule.

235. The method of claim 234 wherein said modulation is *in vivo*.

236. A pharmaceutical composition containing any of the polypeptides in claim 186 in a pharmaceutically acceptable carrier.

237. A pharmaceutically acceptable composition containing any of the nucleic acid molecules of claims 193-195 in a pharmaceutically acceptable carrier.

238. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 193-195.

239. A nonhuman transgenic animal wherein one or more cells of said animal contains any of the nucleic acid sequences of claims 193-195, wherein said cell expresses any of the polypeptides of claim 186.

240. A method for producing a transgenic animal according to claim 239, said method comprising introducing any of the nucleic acid sequences of claims 193-195 into a cell, wherein said cell is present in said animal or gives rise to said animal.

5 241. An agent identified by any of the methods of claims 208-227.

242. An agent identified by any of the methods of claims 229-233.

10

Handwritten notes: "A1" and "A1" with a large curved line.

Sequence length 2175

CACGGTCCGCAAATTTCTGATTCCTTTGAATTAGGATTCCAGATGGGGGCTCATTCTACAGCCCCAACATTCTCT

ATAGCCGTTATCACTGCCATCACCACTGCCACCAGCATCTCTTGCAGATTCCACCCCTGCTCCCCAGAGACTTCTCTGC

TTTGAAAGTGAGCAGAAAGGAAGCTCTCAGAAAAATCTCTAGTGGTGGCTGCCGTCGCTCCAGACAATCGGAATCCTGC

M G W L F L K V L L A G V S F S G	17
CTTCACCACC ATG GGC TGG CTT TTT CTA AAG GTT TTG TTG GCG GGA GTG AGT TTC TCA GGA	51
F L Y P L V D F C I S G K T R G Q K P N	37
TTT CTT TAT CCT CTT GTG GAT TTT TGC ATC AGT GGG AAA ACA AGA GGA CAG AAG CCA AAC	111
F V I I L A D D M G W G D L G A N W A E	57
TTT GTG ATT ATT TTG GCC GAT GAC ATG GGG TGG GGT GAC CTG GGA GCA AAC TGG GCA GAA	171
T K D T A N L D K M A S E G M R F V D F	77
ACA AAG GAC ACT GCC AAC CTT GAT AAG ATG GCT TCG GAG GGA ATG AGG TTT GTG GAT TTC	231
H A A A S T C S P S R A S L L T G R L G	97
CAT GCA GCT GCC TCC ACC TGC TCA CCC TCC CGG GCT TCC TTG CTC ACC GGC CGG CTT GGC	291
L R N G V T R N F A V T S V G G L P L N	117
CTT CGC AAT GGA GTC ACA CGC AAC TTT GCA GTC ACT TCT GTG GGA GGC CTT CCG CTC AAC	351
E T T L A E V L Q Q A G Y V T G I I G K	137
GAG ACC ACC TTG GCA GAG GTG CTG CAG CAG GCG GGT TAC GTC ACT GGG ATA ATA GGC AAA	411
W H L G H H G S Y H P N F R G F D Y Y F	157
TGG CAT CTT GGA CAC CAC GGC TCT TAT CAC CCC AAC TTC CGT GGT TTT GAT TAC TAC TTT	471
G I P Y S H D M G C T D T P G Y N H P P	177
GGA ATC CCA TAT AGC CAT GAT ATG GGC TGT ACT GAT ACT CCA GGC TAC AAC CAC CCT CCT	531
C P A C P Q G D G P S R N L Q R D C Y T	197
TGT CCA GCG TGT CCA CAG GGT GAT GGA CCA TCA AGG AAC CTT CAA AGA GAC TGT TAC ACT	591
D V A L P L Y E N L N I V E Q P V N L S	217
GAC GTG GCC CTC CCT CTT TAT GAA AAC CTC AAC ATT GTG GAG CAG CCG GTG AAC TTG AGC	651
S L A Q K Y A E K A T Q F I Q R A S T S	237
AGC CTT GCC CAG AAG TAT GCT GAG AAA GCA ACC CAG TTC ATC CAG CGT GCA AGC ACC AGC	711
G R P F L L Y V A L A H M H V P L P V T	257
GGG AGG CCC TTC CTG CTC TAT GTG GCT CTG GCC CAC ATG CAC GTG CCC TTA CCC GTG ACT	771
Q L P A A P R G R S L Y G A G L W E M D	277
CAG CTA CCA GCA GCG CCA CGG GGC AGA AGC CTG TAT GGT GCA GGG CTC TGG GAG ATG GAC	831
S L V G Q I K D K V D H T V K E N T F L	297
AGT CTG GTG GGC CAG ATC AAG GAC AAA GTT GAC CAC ACA GTG AAG GAA AAC ACA TTC CTC	891
W F T G D N G P W A Q K C E L A G S V G	317
TGG TTT ACA GGA GAC AAT GGC CCG TGG GCT CAG AAG TGT GAG CTA GCG GGC AGT GTG GGT	951
P F T G F W Q T R Q G G S P A K Q T T W	337
CCC TTC ACT GGA TTT TGG CAA ACT CGT CAA GGG GGA AGT CCA GCC AAG CAG ACG ACC TGG	1011
E G G H R V P A L A Y W P G R V P V N V	357

FIG 1 (-1-)

GAA GGA GGG CAC CGG GTC CCA GCA CTG GCT TAC TGG CCT GGC AGA GTT CCA GTT AAT GTC	1071
T S T A L L S V L D I F P T V V A L A Q	377
ACC AGC ACT GCC TTG TTA AGC GTG CTG GAC ATT TTT CCA ACT GTG GTA GCC CTG GCC CAG	1131
A S L P Q G R R F D G V D V S E V L F G	397
GCC AGC TTA CCT CAA GGA CGG CGC TTT GAT GGT GTG GAC GTC TCC GAG GTG CTC TTT GGC	1191
R S Q P G H R V L F H P N S G A A G E F	417
CGG TCA CAG CCT GGG CAC AGG GTG CTG TTC CAC CCC AAC AGC GGG GCA GCT GGA GAG TTT	1251
G A L Q T V R L E R Y K A F Y I T G G A	437
GGA GCC CTG CAG ACT GTC CGC CTG GAG CGT TAC AAG GCC TTC TAC ATT ACC GGT GGA GCC	1311
R A C D G S T G P E L Q H K F P L I F N	457
AGG GCG TGT GAT GGG AGC ACG GGG CCT GAG CTG CAG CAT AAG TTT CCT CTG ATT TTC AAC	1371
L E D D T A E A V P L E R G G A E Y Q A	477
CTG GAA GAC GAT ACC GCA GAA GCT GTG CCC CTA GAA AGA GGT GGT GCG GAG TAC CAG GCT	1431
V L P E V R K V L A D V L Q D I A N D N	497
GTG CTG CCC GAG GTC AGA AAG GTT CTT GCA GAC GTC CTC CAA GAC ATT GCC AAC GAC AAC	1491
I S S A D Y T Q D P S V T P C C N P Y Q	517
ATC TCC AGC GCA GAT TAC ACT CAG GAC CCT TCA GTA ACT CCC TGC TGT AAT CCC TAC CAA	1551
I A C R C Q A A *	526
ATT GCC TGC CGC TGT CAA GCC GCA TAA	1578
CAGACCAATTTTATTCCACGAGGAGGAGTACCTGGAAATTAGGCAAGTTTGCTTCCAAATTTTCATTTTACCCCTCTTT	
ACAAACACACGCTTTAGTTTAGTCTTGAGTTTAGTTTGGAGTTAGCCTTGCATATCCCTTCTGTATCCTGTCCCTCC	
TCCACGCCGACCCGAGAGCAGCTGAGCTGCGCTGGCTCTGGGCAGGGAGTGTGCCTTAATGGGAAGCACACGGGCTTTG	
GAGTCAGGCACAGGTGCCAGCTCCAGCTTTTGAACCTGGGCAATTGTTTAACCTAACCTGCAAGTTGATTTTGAGGGTT	
AAATAAAGGCATACATGAAAAAAAAAAAAAAAAA	




FIG 2

Prosite Pattern Matches :

Prosite version: Release 12.2 of February 1995

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 117 NETT 120

Query: 215 NLSS 218

Query: 356 NVTG 359

Query: 497 NISS 500

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 28 SGK 30

Query: 93 TGR 95

Query: 237 SGR 239

Query: 290 TVK 292

Query: 422 TVR 424

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 120 TLAE 123

Query: 290 TVKE 293

Query: 335 TTWE 338

Query: 364 SVLD 367

Query: 444 TGPE 447

Query: 499 SSAD 502

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 12 GVSFSG 17

Query: 33 GQKPNF 38

Query: 52 GANWAE 57

Query: 97 GLRNGV 102

Query: 113 GLPLNE 118

Query: 158 GIPYSH 163

Query: 328 GGSPAK 333

Query: 388 GVDVSE 393

Query: 418 GALQTV 423

Query: 435 GGARAC 440

>PS00009|PDOC00009|AMIDATION Amidation site.

Query: 382 QGRR 385

>PS00142|PDOC00117|SULFATASE_2 Sulfatases signature 2.

Query: 129 GYVTGIIGRW 138

Input file Fbh23553f1.seq; Output File 23553.trans
Sequence length 4321

CCACGCGTCCGGCTAATGAATCTTGGGGCGGTGTGGGGCGGGGGCGCTTGATCGGCAACTAGGAAACCCAGGGGC
AGAGGCCAGGAGCGAGGGCAGCGAGGATCAGAGGCCAGGCCTTCCCGGCTGCCGGCGCTCCTCGGAGGTGAGGGCAGAT
GAGGAACATGACTCTCCCCCTTCGGAGGAGGAAGGAAGTCCCGCTGCCACCTTATCTCTGCTCCTCTGCTCCTCCCTG
TTCCAGAGCTTTTCTCTAGAGAAGATTTTGAAGGCGGCTTTTGTGCTGAAGGCCACCCACCATCATCTAAAGAAGAT
AAACTTGGCAAATGACATGCAGGTTCTTCAAGGCAGAATAATTGCAGAAAATCTTCAAAGGACCTATCTGCAGATGTT
CTGAATACCTCTGAGAATAGAGATTGATTATTCAACCAGGATACCTAATTCAAGAACTCCAGAAATCAGGAGACGGAGA

M K Y S C C A L V L A 11
CATTTTGTTCAGTTTTCGAACATTGGACCAATACA ATG AAG TAT TCT TGC TGT GCT CTG GTT TTG GCT 33

V L G T E L L G S L C S T V R S P R F R 31
GTC CTG GGC ACA GAA TTG CTG GGA AGC CTC TGT TCG ACT GTC AGA TCC CCG AGG TTC AGA 93

G R I Q Q E R K N I R P N I I L V L T D 51
GGA CGG ATA CAG CAG GAA CGA AAA AAC ATC CGA CCC AAC ATT ATT CTT GTG CTT ACC GAT 153

D Q D V E L G S L Q V M N K T R K I M E 71
GAT CAA GAT GTG GAG CTG GGG TCC CTG CAA GTC ATG AAC AAA ACG AGA AAG ATT ATG GAA 213

H G G A T F I N A F V T T P M C C P S R 91
CAT GGG GGG GCC ACC TTC ATC AAT GCC TTT GTG ACT ACA CCC ATG TGC TGC CCG TCA CGG 273

S S M L T G K Y V H N H N V Y T N N E N 111
TCC TCC ATG CTC ACC GGG AAG TAT GTG CAC AAT CAC AAT GTC TAC ACC AAC AAC GAG AAC 333

C S S P S W Q A M H E P R T F A V Y L N 131
TGC TCT TCC CCC TCG TGG CAG GCC ATG CAT GAG CCT CGG ACT TTT GCT GTA TAT CTT AAC 393

N T G Y R T A F F G K Y L N E Y N G S Y 151
AAC ACT GGC TAC AGA ACA GCC TTT TTT GGA AAA TAC CTC AAT GAA TAT AAT GGC AGC TAC 453

I P P G W R E W L G L I K N S R F Y N Y 171
ATC CCC CCT GGG TGG CGA GAA TGG CTT GGA TTA ATC AAG AAT TCT CGC TTC TAT AAT TAC 513

T V C R N G I K E K H G F D Y A K D Y F 191
ACT GTT TGT CGC AAT GGC ATC AAA GAA AAG CAT GGA TTT GAT TAT GCA AAG GAC TAC TTC 573

T D L I T N E S I N Y F K M S K R M Y P 211
ACA GAC TTA ATC ACT AAC GAG AGC ATT AAT TAC TTC AAA ATG TCT AAG AGA ATG TAT CCC 633

H R P V M M V I S H A A P H G P E D S A 231
CAT AGG CCC GTT ATG ATG GTG ATC AGC CAC GCT GCG CCC CAC GGC CCC GAG GAC TCA GCC 693

P Q F S K L Y P N A S Q H I T P S Y N Y 251
CCA CAG TTT TCT AAA CTG TAC CCC AAT GCT TCC CAA CAC ATA ACT CCT AGT TAT AAC TAT 753

A P N M D K H W I M Q Y T G P M L P I H 271
GCA CCA AAT ATG GAT AAA CAC TGG ATT ATG CAG TAC ACA GGA CCA ATG CTG CCC ATC CAC 813

M E F T N I L Q R K R L Q T L M S V D D 291
ATG GAA TTT ACA AAC ATT CTA CAG CGC AAA AGG CTC CAG ACT TTG ATG TCA GTG GAT GAT 873

S V E R L Y N M L V E T G E L E N T Y I 311

TCT	GTG	GAG	AGG	CTG	TAT	AAC	ATG	CTC	GTG	GAG	ACG	GGG	GAG	CTG	GAG	AAT	ACT	TAC	ATC	933
I	Y	T	A	D	H	G	Y	H	I	G	Q	F	G	L	V	K	G	K	S	331
ATT	TAC	ACC	GCC	GAC	CAT	GGT	TAC	CAT	ATT	GGG	CAG	TTT	GGA	CTG	GTC	AAG	GGG	AAA	TCC	993
M	P	Y	D	F	D	I	R	V	P	F	F	I	R	G	P	S	V	E	P	351
ATG	CCA	TAT	GAC	TTT	GAT	ATT	CGT	GTG	CCT	TTT	TTT	ATT	CGT	GGT	CCA	AGT	GTA	GAA	CCA	1053
G	S	I	V	P	Q	I	V	L	N	I	D	L	A	P	T	I	L	D	I	371
GGA	TCA	ATA	GTC	CCA	CAG	ATC	GTT	CTC	AAC	ATT	GAC	TTG	GCC	CCC	ACG	ATC	CTG	GAT	ATT	1113
A	G	L	D	T	P	P	D	V	D	G	K	S	V	L	K	L	L	D	P	391
GCT	GGG	CTC	GAC	ACA	CCT	CCT	GAT	GTG	GAC	GGC	AAG	TCT	GTC	CTC	AAA	CTT	CTG	GAC	CCA	1173
E	K	P	G	N	R	F	R	T	N	K	K	A	K	I	W	R	D	T	F	411
GAA	AAG	CCA	GGT	AAC	AGG	TTT	CGA	ACA	AAC	AAG	AAG	GCC	AAA	ATT	TGG	CGT	GAT	ACA	TTC	1233
L	V	E	R	G	K	F	L	R	K	K	E	E	S	S	K	N	I	Q	Q	431
CTA	GTG	GAA	AGA	GGC	AAA	TTT	CTA	CGT	AAG	AAG	GAA	GAA	TCC	AGC	AAG	AAT	ATC	CAA	CAG	1293
S	N	H	L	P	K	Y	E	R	V	K	E	L	C	Q	Q	A	R	Y	Q	451
TCA	AAT	CAC	TTG	CCC	AAA	TAT	GAA	CGG	GTC	AAA	GAA	CTA	TGC	CAG	CAG	GCC	AGG	TAC	CAG	1353
T	A	C	E	Q	P	G	Q	K	W	Q	C	I	E	D	T	S	G	K	L	471
ACA	GCC	TGT	GAA	CAA	CCG	GGG	CAG	AAG	TGG	CAA	TGC	ATT	GAG	GAT	ACA	TCT	GGC	AAG	CTT	1413
R	I	H	K	C	K	G	P	S	D	L	L	T	V	R	Q	S	T	R	N	491
CGA	ATT	CAC	AAG	TGT	AAA	GGA	CCC	AGT	GAC	CTG	CTC	ACA	GTC	CGG	CAG	AGC	ACG	CGG	AAC	1473
L	Y	A	R	G	F	H	D	K	D	K	E	C	S	C	R	E	S	G	Y	511
CTC	TAC	GCT	CGC	GGC	TTC	CAT	GAC	AAA	GAC	AAA	GAG	TGC	AGT	TGT	AGG	GAG	TCT	GGT	TAC	1533
R	A	S	R	S	Q	R	K	S	Q	R	Q	F	L	R	N	Q	G	T	P	531
CGT	GCC	AGC	AGA	AGC	CAA	AGA	AAG	AGT	CAA	CGG	CAA	TTC	TTG	AGA	AAC	CAG	GGG	ACT	CCA	1593
K	Y	K	P	R	F	V	H	T	R	Q	T	R	S	L	S	V	E	F	E	551
AAG	TAC	AAG	CCC	AGA	TTT	GTC	CAT	ACT	CGG	CAG	ACA	CGT	TCC	TTG	TCC	GTC	GAA	TTT	GAA	1653
G	E	I	Y	D	I	N	L	E	E	E	E	E	L	Q	V	L	Q	P	R	571
GGT	GAA	ATA	TAT	GAC	ATA	AAT	CTG	GAA	GAA	GAA	GAA	GAA	TTG	CAA	GTG	TTG	CAA	CCA	AGA	1713
N	I	A	K	R	H	D	E	G	H	K	G	P	R	D	L	Q	A	S	S	591
AAC	ATT	GCT	AAG	CGT	CAT	GAT	GAA	GGC	CAC	AAG	GGG	CCA	AGA	GAT	CTC	CAG	GCT	TCC	AGT	1773
G	G	N	R	G	R	M	L	A	D	S	S	N	A	V	G	P	P	T	T	611
GGT	GGC	AAC	AGG	GGC	AGG	ATG	CTG	GCA	GAT	AGC	AGC	AAC	GCC	GTG	GGC	CCA	CCT	ACC	ACT	1833
V	R	V	T	H	K	C	F	I	L	P	N	D	S	I	H	C	E	R	E	631
GTC	CGA	GTG	ACA	CAC	AAG	TGT	TTT	ATT	CTT	CCC	AAT	GAC	TCT	ATC	CAT	TGT	GAG	AGA	GAA	1893
L	Y	Q	S	A	R	A	W	K	D	H	K	A	Y	I	D	K	E	I	E	651
CTG	TAC	CAA	TCG	GCC	AGA	GCG	TGG	AAG	GAC	CAT	AAG	GCA	TAC	ATT	GAC	AAA	GAG	ATT	GAA	1953
A	L	Q	D	K	I	K	N	L	R	E	V	R	G	H	L	K	R	R	K	671
GCT	CTG	CAA	GAT	AAA	ATT	AAG	AAT	TTA	AGA	GAA	GTG	AGA	GGA	CAT	CTG	AAG	AGA	AGG	AAG	2013
P	E	E	C	S	C	S	K	Q	S	Y	Y	N	K	E	K	G	V	K	K	691
CCT	GAG	GAA	TGT	AGC	TGC	AGT	AAA	CAA	AGC	TAT	TAC	AAT	AAA	GAG	AAA	GGT	GTA	AAA	AAG	2073
Q	E	K	L	K	S	H	L	H	P	F	K	E	A	A	Q	E	V	D	S	711
CAA	GAG	AAA	TTA	AAG	AGC	CAT	CTT	CAC	CCA	TTC	AAG	GAG	GCT	GCT	CAG	GAA	GTA	GAT	AGC	2133

FIG 5 (-2-)

K	L	Q	L	F	K	E	N	N	R	R	R	K	K	E	R	K	E	K	R	731
AAA	CTG	CAA	CTT	TTC	AAG	GAG	AAC	AAC	CGT	AGG	AGG	AAG	AAG	GAG	AGG	AAG	GAG	AAG	AGA	2193
R	Q	R	K	G	E	E	C	S	L	P	G	L	T	C	F	T	H	D	N	751
CGG	CAG	AGG	AAG	GGG	GAA	GAG	TGC	AGC	CTG	CCT	GGC	CTC	ACT	TGC	TTC	ACG	CAT	GAC	AAC	2253
N	H	W	Q	T	A	P	F	W	N	L	G	S	F	C	A	C	T	S	S	771
AAC	CAC	TGG	CAG	ACA	GCC	CCG	TTC	TGG	AAC	CTG	GGA	TCT	TTC	TGT	GCT	TGC	ACG	AGT	TCT	2313
N	N	N	T	Y	W	C	L	R	T	V	N	E	T	H	N	F	L	F	C	791
AAC	AAT	AAC	ACC	TAC	TGG	TGT	TTG	CGT	ACA	GTT	AAT	GAG	ACG	CAT	AAT	TTT	CTT	TTC	TGT	2373
E	F	A	T	G	F	L	E	Y	F	D	M	N	T	D	P	Y	Q	L	T	811
GAG	TTT	GCT	ACT	GGC	TTT	TTG	GAG	TAT	TTT	GAT	ATG	AAT	ACA	GAT	CCT	TAT	CAG	CTC	ACA	2433
N	T	V	H	T	V	E	R	G	I	L	N	Q	L	H	V	Q	L	M	E	831
AAT	ACA	GTG	CAC	ACG	GTA	GAA	CGA	GGC	ATT	TTG	AAT	CAG	CTA	CAC	GTA	CAA	CTA	ATG	GAG	2493
L	R	S	C	Q	G	Y	K	Q	C	N	P	R	P	K	N	L	D	V	G	851
CTC	AGA	AGC	TGT	CAA	GGA	TAT	AAG	CAG	TGC	AAC	CCA	AGA	CCT	AAG	AAT	CTT	GAT	GTT	GGA	2553
N	K	D	G	G	S	Y	D	L	H	R	G	Q	L	W	D	G	W	E	G	871
AAT	AAA	GAT	GGA	GGA	AGC	TAT	GAC	CTA	CAC	AGA	GGA	CAG	TTA	TGG	GAT	GGA	TGG	GAA	GGT	2613
*																				872
TAA																				2616

TCAGCCCCGTCTCACTGCAGACATCAACTGGCAAGGCCTAGAGGAGCTACACAGTGTGAATGAAAACATCTATGAGTAC
AGACAAAACCTACAGACTTAGTCTGGTGGACTGGACTAATTACTTGAAGGATTAGATAGAGTATTTGCACTGCTGAAGA
GTCACATATGAGCAAAATAAAACAAATAAGACTCAAAGTCAAGTGACGGGTTCTTGGTTGTCTCTGCTGAGCACGC
TGTGTCAATGGAGATGGCCTCTGCTGACTCAGATGAAGACCCAGGCATAAGGTTGGGAAAAACACCTCATTTGACCTTG
CCAGCTGACCTTCAAACCCCTGCATTTGAACCGACCAACATTAAGTCCAGAGAGTAAACTTGAATGGAATAACGACATTC
CAGAAGTTAATCATTTGAATTCTGAACACTGGAGAAAAACCGAAAAATGGACGGGGCATGAAGAGACTAATCATCTGGA
AACCGATTTCAGTGGCGATGGCATGACAGAGCTAGAGCTCGGGCCCGAGCCCGAGGCTGCAGCCCATTCGCAGGCACCCG
AAAGAACTTCCCCAGTATGGTGGTCCCTGGAAAAGGACATTTTGAAGATCAACTATATCTTCTGTGCATTCGGATGGAA
TTTCAGTTCATCAGATGTTACCATGGCCACCGCAGAACACCGAAGTAATCCAGCATAGCGGGGAAGATGTTGACCAA
GGTGGAGAAGAATCAAGAAAAGGAGAAGTACAGCACCTAGAAGGCAGCGCTCCTCTTCACTCTCTCTGATTAGATG
AAACTGTTACCTTACCTAAACACAGTATTTCTTTTAACTTTTATTTGTAAACTAATAAAGGKAATCACAGCCACC
AACATTCCAAGCTACCTGGGTACCTTTGTGCACTAGAAGCTAGTGAGCATGTGAGCAAGCGGTGTGCACACGGAGACT
CATGTTATAATTTACTATCTGCCAAGGAGTAGAAAGAAAGGCTGGGGATATTTGGGTGGCTTTGGKTTTGATTTT
GCTTGGTTGGTTGGTTTGKACTAAAACAGTATTATCTTTGAATATCGTAGGGACATAARKWWWMMWKIWTTCMAW
YMRKAKGSYWRRAWKGGGSTYTYTSKKRKSTMWAMWYKWSMCCYSKKRWAWTYWYMMYWCYKYTSSSTGRYKRN
KTAATGAAGTT

Analysis of 23553 (871 aa)




FIG 6




FIG 7

Prosite Pattern Matches for 23553

Prosite version: Release 12.2 of February 1995

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 64	NKTR	67
Query: 111	NCSS	114
Query: 131	NNTG	134
Query: 148	NGSY	151
Query: 170	NYTV	173
Query: 197	NESI	200
Query: 240	NASQ	243
Query: 623	NDSI	626
Query: 773	NNTY	776
Query: 783	NETH	786

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 24	TVR	26
Query: 27	SPR	29
Query: 66	TRK	68
Query: 96	TGK	98
Query: 206	SKR	208
Query: 400	TNK	402
Query: 425	SSK	427
Query: 468	SGK	470
Query: 484	TVR	486
Query: 488	STR	490
Query: 505	SCR	507
Query: 516	SQR	518
Query: 520	SQR	522
Query: 530	TPK	532
Query: 611	TVR	613
Query: 615	THK	617
Query: 635	SAR	637

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 107	TNNE	110
Query: 288	SVDD	291
Query: 367	TILO	370
Query: 376	TPPD	379
Query: 452	TACE	455
Query: 505	SCRE	508
Query: 781	TVNE	784

>PS00007|PDOC00007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

Query: 637 RAWKDHKAY 645

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 19 GSLCST 24

Query: 161 GLIKNS 166

Query: 325 GLVKGK 330

Query: 592 GGNRGR 597

Query: 763 GSFCAC 768

Query: 851 GNKDGG 856

>PS00523|PDOC00117|SULFATASE_1 Sulfatases signature 1.

Query: 85 PMCCPSRSSMLTG 97




FIG 9

Input file Fbh25278FL1.seq; Output File 25278.trans
Sequence length 2940

CCACGCGTCGCCACGCGTCCGGCTGCCACGCCGCTCTCAGGCTGGCCGGGCTGAGCCGGGAAGAGGGAGCAAAGG
CGGCGCAGGGCTCGCGTTAGGCAGCGGGAGGCAGCTCGCGCGGGGCTGACCTCCCCAGAGCGCCCGCTGCGGCCGA
GCAGATCCGGCCAGCCGTCGGGAGCCAGTCCCGGACCAGACACTGGACCGTCCCCGGGGGGCGCTGAACCTCCCTCGC
AGCATCCGAGCCGGCGGGCCGGTGGTGGCCCTGGGCGCGCGAGGTGGTGAGGCCCCAGGAGCCCGGCGCGCCGGGACA

M	H	T	L	T	G	F	S	L	V	S	L	L	S	F	15
CGCGGGCCGGCTTGGCG	ATG	CAC	ACC	CTC	ACT	GGC	TTC	TCT	CTG	GTC	AGC	CTG	CTC	AGC	45
G	Y	L	S	W	D	W	A	K	P	S	F	V	A	D	35
GGC	TAC	CTG	TCC	TGG	GAC	TGG	GCC	AAG	CCG	AGC	TTC	GTG	GCC	GAC	105
G	E	Q	P	S	A	A	P	P	Q	P	P	H	I	I	55
GGC	GAG	CAG	CCC	TCG	GCC	GCT	CCG	CCC	CAG	CCT	CCC	CAC	ATC	ATC	165
D	Q	G	Y	H	D	V	G	Y	H	G	S	D	I	E	75
GAC	CAA	GGC	TAC	CAC	GAC	GTG	GGC	TAC	CAT	GGT	TCA	GAT	ATC	GAG	225
R	L	A	A	K	G	V	K	L	E	N	Y	Y	I	Q	95
AGG	CTG	GCG	GCC	AAG	GGG	GTC	AAG	TTG	GAG	AAT	TAT	TAC	ATC	CAG	285
S	R	S	Q	L	L	T	G	R	Y	Q	I	H	T	G	115
TCG	CGG	AGC	CAG	CTC	CTC	ACT	GGC	AGG	TAC	CAG	ATC	CAC	ACA	GGA	345
I	R	P	Q	Q	P	N	C	L	P	L	D	Q	V	T	135
ATC	CGC	CCA	CAG	CAG	CCC	AAC	TGC	CTG	CCC	CTG	GAC	CAG	GTG	ACA	405
Q	E	A	G	Y	S	T	H	M	V	G	K	W	H	L	155
CAG	GAG	GCA	GGT	TAT	TCC	ACC	CAT	ATG	GTG	GGC	AAG	TGG	CAC	CTG	465
E	C	L	P	T	R	R	G	F	D	T	F	L	G	S	175
GAG	TGT	CTG	CCC	ACC	CGT	CGG	GGC	TTC	GAC	ACC	TTC	CTG	GGC	TCG	525
D	Y	Y	T	Y	D	N	C	D	G	P	G	V	C	G	195
GAC	TAT	TAC	ACC	TAT	GAC	AAC	TGT	GAT	GGC	CCA	GGC	GTG	TGC	GGC	585
G	E	N	V	A	W	G	L	S	G	Q	Y	S	T	M	215
GGT	GAG	AAT	GTG	GCC	TGG	GGG	CTC	AGC	GGC	CAG	TAC	TCC	ACT	ATG	645
A	S	H	I	L	A	S	H	S	P	Q	R	P	L	F	235
GCC	AGC	CAT	ATC	CTG	GCC	AGC	CAC	AGC	CCT	CAG	CGT	CCC	CTC	TTC	705
Q	A	V	H	T	P	L	Q	S	P	R	E	Y	L	Y	255
CAG	GCA	GTA	CAC	ACA	CCC	CTG	CAG	TCC	CCT	CGT	GAG	TAC	CTG	TAC	765
G	N	V	A	R	R	K	Y	A	A	M	V	T	C	M	275
GGC	AAT	GTG	GCC	CGG	CGG	AAG	TAC	GCG	GCC	ATG	GTG	ACC	TGC	ATG	825
N	I	T	W	A	L	K	R	Y	G	F	Y	N	N	S	295
AAC	ATC	ACC	TGG	GCC	CTC	AAG	CGC	TAC	GGT	TTC	TAC	AAC	AAC	AGT	885
S	D	N	G	G	Q	T	F	S	G	G	S	N	W	P	315
AGT	GAC	AAT	GGT	GGC	CAG	ACT	TTC	TCG	GGG	GGC	AGC	AAC	TGG	CCG	945
G	T	Y	W	E	G	G	V	R	G	L	G	F	V	H	335
GGC	ACT	TAT	TGG	GAA	GGT	GGC	GTG	CGG	GGC	CTA	GGC	TTT	GTC	CAC	1005

R K Q R T S R A L M H I T D W Y P T L V	355
CCA AAG CAA CGG ACA AGC CGG GCA CTG ATG CAC ATC ACT GAC TGG TAC CCG ACC CTG GTG	1065
G L A G G T T S A A D G L D G Y D V W P	375
GGT CTG GCA GGT GGT ACC ACC TCA GCA GCC GAT GGG CTA GAT GGC TAC GAC GTG TGG CCG	1125
A I S E G R A S P R T E I L H N I D P L	395
GCC ATC AGC GAG GGC CGG GCC TCA CCA CGC ACG GAG ATC CTG CAC AAC ATT GAC CCA CTC	1185
Y N H A Q H G S L E G G F G I W N T A V	415
TAC AAC CAT GCC CAG CAT GGC TCC CTG GAG GGC GGC TTT GGC ATC TGG AAC ACC GCC GTG	1245
Q A A I R V G E W K L L T G D P G Y G D	435
CAG GCT GCC ATC CGC GTG GGT GAG TGG AAG CTG CTG ACA GGA GAC CCC GGC TAT GGC GAT	1305
W I P P Q T L A T F P G S W W N L E R M	455
TGG ATC CCA CCG CAG ACA CTG GCC ACC TTC CCG GGT AGC TGG TGG AAC CTG GAA CGA ATG	1365
A S V R Q A V W L F N I S A D P Y E R E	475
GCC AGT GTC CGC CAG GCC GTG TGG CTC TTC AAC ATC AGT GCT GAC CCT TAT GAA CGG GAG	1425
D L A G Q R P D V V R T L L A R L A E Y	495
GAC CTG GCT GGC CAG CGG OCT GAT GTG GTC CGC ACC CTG CTG GCT CGC CTG GCC GAA TAT	1485
N R T A I P V R Y P A E N P R A H P D F	515
AAC CGC ACA GCC ATC CCG GTA CGC TAC CCA GCT GAG AAC CCC CGG GCT CAT CCT GAC TTT	1545
N G G A W G P W A S D E E E E E E E G R	535
AAT GGG GGT GCT TGG GGG CCC TGG GCC AGT GAT GAG GAA GAG GAG GAA GAG GAA GGG AGG	1605
A R S F S R G R R K K K C K I C K L R S	555
GCT CGA AGC TTC TCC CGG GGT GGT CGC AAG AAA AAA TGC AAG ATT TGC AAG CTT CGA TCC	1665
F F R K L N T R L M S Q R I *	570
TTT TTC CGT AAA CTC AAC ACC AGG CTA ATG TCC CAA CGG ATC TGA	1710

TGGTGGGGAGGGAGAAAACGTCTTTAGAGGATCTTCCCCACTCCGGCTTGGCCCTGCTGTCTCTCAGGGAGAAGCCT
 GTACATCTCCATCTACAGGGAGTTGGAGGGTGTAGAGTCCCTTGGTTGAACAGGGTAGGGAGCCTGGATAGGAGTGGG
 TGGGAATAAACAGACTGGGATGCCGTGTGTCTCAGTCTGCTCTCCTCAAGGACTTGCTCTGTGACCTCAGGTGACCCAC
 ATGAGCTTTTAGCCTCAGTTTCTCATCTGTAAATGAGCTCTAATGACTTTGTGACTCTTTGGTGTGGCCCTGGAGCC
 TGGGGCACGGTGGAGTTCTTGGCCGGCCTTGCCACTTGACAACCTCTTAAGGCTTCCCCCTTAACAAGGGATCCCTG
 TGGTGGTGTTTGGGAGTTGCCGTGAGGCAACTCCAAGCCTGGCCCCAGCTGAAGCATGGCAATCTGGCTGCTCTCTAC
 AGGGACCCCCAAGCGCTGTGGGTGGAGGGCAGGGGTGGGGGGGTGACCTTCTTGGGTCTTCACATGGCCTAGGCCAG
 TCCTCCGGTCAGACTGGTGTGAGGCACCGTGGTGCAAAATTCCTCTTCTGGCCCCCTCCAGTACCCAGAGAACTGGCTG
 GGCCATTAACTGCTGCAGCACCAAGGTGGTAGAAAGAGCTGTGAAGAGCCCCAAACCAGTACCAGGACACCTGGGTT
 CTCTGTGACCTGGGGCACAGTTCTTGGCCTCTAGGCCTTGATTTCGCCACCTGCAAGTGGGGATGCCAGCCCTGGCTC
 TGCCCTCTTCATGAGGCTCTGGAAGACTGGCCAAGGTGTGGAGGAGCTGTGAACTTGATTAAAGTGTGTAACATGG
 AAAAAAAAAAAAAAAAAAAGGGCGG

Analysis of 25278 (569 aa)




FIG- 11




FIG 12

Prosite Pattern Matches for 2578

Prosite version: Release 12.2 of February 1998

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 276 NITW 279
Query: 288 MMSV 291
Query: 466 NISA 469
Query: 496 NRTA 499

>PS00004|PDOC00004|CAMP_PHOSPHO_SITE cAMP- and cGMP-dependent protein kinase phosphorylation site.

Query: 314 RRGY 317

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 102 TGR 104
Query: 160 TRR 162
Query: 244 SPR 246
Query: 340 TSR 342
Query: 383 SPR 385
Query: 457 SVR 459
Query: 566 SQR 568

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 67 SDIE 70
Query: 244 SPRE 247
Query: 268 TCMD 271
Query: 317 TYWE 320
Query: 363 SAAD 366
Query: 525 SDEE 528

>PS00007|PDOC00007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

Query: 134 KLQEAGY 140

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 110 GLQHSI 115
Query: 169 GSLTGN 174
Query: 205 GQYSTH 210
Query: 300 GQTFSG 305
Query: 321 GGVRLG 326
Query: 356 GLAGGT 361
Query: 402 GSLEGG 407
Query: 409 GIWNTA 414
Query: 447 GSWWNL 452

>PS00009|PDOC00009|AMIDATION Amidation site.

Query: 312 RGRK 315
Query: 541 RGRR 544

>PS00142|PDOC00117|SULFATASE_2 Sulfatases signature 2.

Query: 139 GYSTHVGKW 148

>PS00523|PDOC00117|SULFATASE_1 Sulfatases signature 1.

Query: 91 PICTPSRSQLLTG 103




FIG 14

Input file 26212cons; Output File 26212pat
Sequence length 2266

CACGCGTCCGCCCCACGCGTCCGTGGAGATATTAACCTTTTTCTTTTTTTTTTCCTTGGTGGAAAGCTGCTCTAGGGAGG
GGGGAGGAGGAGGAGAAAAGTGAAATGTGCTGGAGAAGAGCGAGCCCTCCTTGTCTTCCGGAGTCCCATCCATTAAGCC
ATCACTTCTGGAAGATTAAAGTTGTGCGACATGGTGACAGCTGAGAGGAGAGGAGGATTTCCTTGCCAGGTGGAGAGTCT
TCACCGTCTGTTGGGTGCATGTGTGCGCCCGCASC GCGCGGGGCGCGTGGTTCTCCGCGTGGAGTCTCACCTGGGACC

M	A	P	R	G	C	A	G	H	P	P	P	P	S	P	Q	A	C	18
ATG	GCT	CCC	AGG	GGC	TGT	GCG	GGG	CAT	CCG	CCT	CCG	CCT	TCT	CCA	CAG	GCC	TGT	54
V	C	P	G	K	M	L	A	M	G	A	L	A	G	F	W	I	L	38
GTC	TGT	CCT	GGA	AAG	ATG	CTA	GCA	ATG	GGG	GCG	CTG	GCA	GGA	TTC	TGG	ATC	CTC	114
L	T	Y	G	Y	L	S	W	G	Q	A	L	E	E	E	E	E	G	58
CTC	ACT	TAT	GGT	TAC	CTG	TCC	TGG	GGC	CAG	GCC	TTA	GAA	GAG	GAG	GAA	GAA	GGG	174
L	A	Q	A	G	E	K	L	E	P	S	T	T	S	T	S	Q	P	78
CTA	GCT	CAA	GCT	GGA	GAG	AAA	CTA	GAG	CCC	AGC	ACA	ACT	TCC	ACC	TCC	CAG	CCC	234
I	F	I	L	A	D	D	Q	G	F	R	D	V	G	Y	H	G	S	98
ATT	TTC	ATC	CTA	GCG	GAT	GAT	CAG	GGA	TTT	AGA	GAT	GTG	GGT	TAC	CAC	GGA	TCT	294
T	P	T	L	D	K	L	A	A	E	G	V	K	L	E	N	Y	Y	118
A	ACA	CCT	ACT	CTT	GAC	AAG	CTC	GCT	GCC	GAA	GGA	GTT	AAA	CTG	GAG	AAC	TAC	354
P	I	C	T	P	S	R	S	Q	F	I	T	G	K	Y	Q	I	H	138
G	CCT	ATT	TGC	ACA	CCA	TCC	AGG	AGT	CAG	TTT	ATT	ACT	GGA	AAG	TAT	CAG	ATA	414
L	Q	H	S	I	I	R	P	T	Q	P	N	C	L	P	L	D	N	158
A	CTT	CAA	CAT	TCT	ATC	ATA	AGA	CCT	ACC	CAA	CCC	AAC	TGT	TTA	CCT	CTG	GAC	474
L	P	Q	K	L	K	E	V	G	Y	S	T	H	M	V	G	K	W	178
C	CTA	CCT	CAG	AAA	CTG	AAG	GAG	GTT	GGA	TAT	TCA	ACG	CAT	ATG	GTC	GGA	AAA	534
G	F	Y	R	K	E	C	M	P	T	R	R	G	F	D	T	F	F	198
G	GGT	TTT	TAC	AGA	AAA	GAA	TGC	ATG	CCC	ACC	AGA	AGA	GGA	TTT	GAT	ACC	TTT	594
L	L	G	S	G	D	Y	Y	T	H	Y	K	C	D	S	P	G	M	218
C	CTT	TTG	GGA	AGT	GGG	GAT	TAC	TAT	ACA	CAC	TAC	AAA	TGT	GAC	AGT	CCT	GGG	654
Y	D	L	Y	E	N	D	N	A	A	W	D	Y	D	N	G	I	Y	238
C	TAT	GAC	TTG	TAT	GAA	AAC	GAC	AAT	GCT	GCC	TGG	GAC	TAT	GAC	AAT	GGC	ATA	714
T	Q	M	Y	T	Q	R	V	Q	Q	I	L	A	S	H	N	P	T	258
ACA	CAG	ATG	TAC	ACT	CAG	AGA	GTA	CAG	CAA	ATC	TTA	GCT	TCC	CAT	AAC	CCC	ACA	774
I	F	L	Y	I	A	Y	Q	A	V	H	S	P	L	Q	A	P	G	278
ATA	TTT	TTA	TAT	ATT	GCC	TAT	CAA	GCT	GTT	CAT	TCA	CCA	CTG	CAA	GCT	CCT	GGC	834
F	E	H	Y	R	S	I	I	N	I	N	R	R	R	Y	A	A	M	298
TTC	GAA	CAC	TAC	CGA	TCC	ATT	ATC	AAC	ATA	AAC	AGG	AGG	AGA	TAT	GCT	GCC	ATG	894
C	L	D	E	A	I	N	N	V	T	L	A	L	K	T	Y	G	F	318
TGC	TTA	GAT	GAA	GCA	ATC	AAC	AAC	GTG	ACA	TTG	GCT	CTA	AAG	ACT	TAT	GGT	TTC	954
N	S	I	I	I	Y	S	S	D	N	G	G	Q	P	T	A	G	G	338
AAC	AGC	ATT	ATC	ATT	TAC	TCT	TCA	GAT	AAT	GGT	GGC	CAG	CCT	ACG	GCA	GGA	GGG	1014
W	P	L	R	G	S	K	G	T	Y	W	E	G	G	I	R	A	V	358
TGG	CCT	CTC	AGA	GGT	AGC	AAA	GGA	ACA	TAT	TGG	GAA	GGA	GGG	ATC	CGG	GCT	GTA	1074


FIG 15 (1)

V	H	S	P	L	L	K	N	K	G	T	V	C	K	E	L	V	H	I	T	378
GTG	CAT	AGC	CCA	CTT	CTG	AAA	AAC	AAG	GGA	ACA	GTG	TGT	AAG	GAA	CTT	GTG	CAC	ATC	ACT	1134
D	W	Y	P	T	L	I	S	L	A	E	G	Q	I	D	E	D	I	Q	L	398
GAC	TGG	TAC	CCC	ACT	CTC	ATT	TCA	CTG	GCT	GAA	GGA	CAG	ATT	GAT	GAG	GAC	ATT	CAA	CTA	1194
D	G	Y	D	I	W	E	T	I	S	E	G	L	R	S	P	R	V	D	I	418
GAT	GGC	TAT	GAT	ATC	TGG	GAG	ACC	ATA	AGT	GAG	GGT	CTT	CGC	TCA	CCC	CGA	GTA	GAT	ATT	1254
L	H	N	I	D	P	I	Y	T	K	A	K	N	G	S	W	A	A	G	Y	438
TTG	CAT	AAC	ATT	GAC	CCC	ATA	TAC	ACC	AAG	GCA	AAA	AAT	GGC	TCC	TGG	GCA	GCA	GGC	TAT	1314
G	I	W	N	T	A	I	Q	S	A	I	R	V	Q	H	W	K	L	L	T	458
GGG	ATC	TGG	AAC	ACT	GCA	ATC	CAG	TCA	GCC	ATC	AGA	GTG	CAG	CAC	TGG	AAA	TTG	CTT	ACA	1374
G	N	P	G	Y	S	D	W	V	P	P	Q	S	F	S	N	L	G	P	N	478
GGA	AAT	CCT	GGC	TAC	AGC	GAC	TGG	GTC	CCC	CCT	CAG	TCT	TTC	AGC	AAC	CTG	GGA	CCG	AAC	1434
R	W	H	N	E	R	I	T	S	S	T	G	K	S	V	W	L	F	N	I	498
CGG	TGG	CAC	AAT	GAA	CGG	ATC	ACC	TCG	TCA	ACT	GGC	AAA	AGT	GTA	TGG	CTT	TTC	AAC	ATC	1494
T	A	D	P	Y	E	R	V	D	L	S	N	R	Y	P	G	I	V	K	K	518
ACA	GCC	GAC	CCA	TAT	GAG	AGG	GTG	GAC	CTA	TCT	AAC	AGG	TAT	CCA	GGA	ATC	GTG	AAG	AAG	1554
L	R	R	L	S	Q	F	N	K	T	A	V	P	V	R	Y	P	P	K		538
CTA	CGG	AGG	CTC	TCA	CAG	TTC	AAC	AAA	ACT	GCA	GTG	CCG	GTC	AGG	TAT	CCC	CCC	AAA		1614
P	R	S	N	P	R	L	N	G	G	V	*									551
CCC	AGA	AGT	AAC	CCT	AGG	CTC	AAT	GGA	GGG	GTC	TAG									1653

ACCATGGTATAGAGAGGAAACCAAGAAAAAGAAGCCAAGCAAAAATCAGGCTGAGAAAAAGCAAAAGAAAAGCAAAA
 AGAAGAAGAAACAGCAGAAAGCAGTCTCAGGTTCAACTTGCCATTGAGGTGTTACTTGTGGATAAGCACAAATATTT
 TTTTGGTTAACTTTAATCAGTTCTTATCTTTTCATCTGTTTCTAGGTAAACCAGCAAATTTGGCTCGATAATATC
 TGGCCTAAGCGTCAGGCTTGTTTTCATGCTGTGCCACCTGGTGCCGAATTC

FIG 15 (2)

Analysis of 26212prot (550 aa)






FIG 17

Prosite Pattern Matches for 26212prot

Prosite version: Release 12.2 of February 1995

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 157 NATL 160

Query: 306 NVTL 309

Query: 318 NNSI 321

Query: 431 NGSW 434

Query: 497 NITA 500

Query: 527 NKTA 530

>PS00004|PDOC00004|CAMP_PHOSPHO_SITE cAMP- and cGMP-dependent protein kinase phosphorylat

Query: 521 RRLS 524

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 131 TGK 133

Query: 189 TRR 191

Query: 243 TQR 245

Query: 413 SPR 415

Query: 489 TGK 491

Query: 509 SNR 511

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 298 SCLD 301

Query: 347 TYWE 350

Query: 386 SLAE 389

Query: 406 TISE 409

>PS00007|PDOC00007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

Query: 163 KLKEVG 169

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 28 GALAGF 33

Query: 56 GALLAQ 61

Query: 139 GLQHSI 144

FIG 18 (C)

Query: 198 GSSLGS 203
Query: 235 GIYSTQ 240
Query: 329 GGQPTA 334
Query: 343 GSKGTY 348
Query: 351 GGIRAV 356
Query: 432 GSWAAG 437
Query: 439 GIWNTA 444

>PS00149|PDOC00117|SULFATASE_2 Sulfatases signature 2.

Query: 168 GYSTHVGKW 177

>PS00523|PDOC00117|SULFATASE_1 Sulfatases signature 1.

Query: 120 PICTPSRSQFITG 132

#3

Attorney's Docket No. 5800-79

PATENT



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: Glucksmann et al.

Appl No.: 09/495,823

Group Art Unit: 1643

Filed: January 31, 2000

For: 22438, 23553, 25278, AND 26212 NOVEL HUMAN SULFATASES

June 5, 2000

Box Missing Parts

Assistant Commissioner for Patents

Washington, DC 20231

**SUBMITTAL OF DECLARATION
UNDER 37 C.F.R. § 1.63**

Sir:

Enclosed is a Declaration and Power of Attorney for the above-identified patent application that has been executed by the named inventors. Also enclosed are a Preliminary Amendment and Information Disclosure Statement.

A check in the amount of \$928 is also enclosed to cover the basic filing fee (\$798) and the surcharge under 37 C.F.R. § 1.16(e) (\$130). Any additional fee or credit may be charged to our Deposit Account No. 16-0605.

Respectfully submitted,

W. Murray Spruill
Registration No. 32,943

ALSTON & BIRD LLP

Post Office Drawer 34009

Charlotte, NC 28234-4009

Tel Raleigh Office (919) 420-2200


Fax Raleigh Office (919) 420-2260

RTA01/2078862v1

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Box Missing Parts, Assistant Commissioner For Patents, Washington, DC 20231, on June 5, 2000.

Nancy Cochran

**DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

22438, 23553, 25278, AND 26212, NOVEL HUMAN SULFATASES,

the specification of which

☐ is attached hereto

OR

☒ was filed on January 31, 2000 as United States Application No. 09/495,823.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or

§ 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign application for patent or inventor's certificate, or of any PCT International application having a filing date before that of the application on which priority is claimed.

			<input type="checkbox"/> Yes <input type="checkbox"/> No
Number	Country	MM/DD/YYYY Filed	Priority Claimed
			<input type="checkbox"/> Yes <input type="checkbox"/> No
Number	Country	MM/DD/YYYY Filed	Priority Claimed

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

Application Number	Filing Date (MM/DD/YYYY)
Application Number	Filing Date (MM/DD/YYYY)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application (37 C.F.R. § 1.63(d)).

Application No.	Filing Date	Status (Patented/Pending/Abandoned)
Application. No.	Filing Date	Status (Patented/Pending/Abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following practitioners:

Mark Boshar, Reg. No. 35,456
Jean M. Silveri, Reg. No. 39,030
Theodore Allen, Reg. No. 41,578
Cynthia Kanik, Reg. No. 37,320
Scott A. Brown, Reg. No. 32,724

and, I also appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:


Customer Number 826

Direct correspondence to the
attention of and telephone calls to:

W. Murray Spruill
Registration No. 32,943
Tel Raleigh Office (919) 420-2200
Fax Raleigh Office (919) 420-2260

Full name of (first/sole) inventor: Maria Alexandra Glucksmann

Inventor's

Signature: X 

Date: X 4-21-00

Residence:

33 Summit Road, Lexington, MA 02173, USA

Citizenship:


AR

Post Office Address:

Same as above

Full name of second inventor: Mark Williamson

Inventor's

Signature: X 

Date: X 4.15.00

Residence:

15 Stonecrest Drive, Saugus, MA 01906, USA

Citizenship:

US

Post Office Address:

Same as above